

Circulation

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Myocardial Gene Transfer and Overexpression of β 2-Adrenergic Receptors Potentiates the Functional Recovery of Unloaded Failing Hearts

Hendrik T. Tevæarai, Andrea D. Eckhart, G. Brant Walton, Janelle R. Keys, Katrina Wilson and Walter J. Koch

Circulation 2002;106;124-129; originally published online Jun 17, 2002;

DOI: 10.1161/01.CIR.0000020220.79105.FD

Circulation is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214
Copyright © 2002 American Heart Association. All rights reserved. Print ISSN: 0009-7322. Online
ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circ.ahajournals.org/cgi/content/full/106/1/124>

Subscriptions: Information about subscribing to *Circulation* is online at
<http://circ.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax: 410-528-8550. E-mail:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/reprints>

Myocardial Gene Transfer and Overexpression of β_2 -Adrenergic Receptors Potentiates the Functional Recovery of Unloaded Failing Hearts

Hendrik T. Tevæarai, MD; Andrea D. Eckhart, PhD; G. Brant Walton, BS; Janelle R. Keys, PhD; Katrina Wilson, BS; Walter J. Koch, PhD

Background—Mechanical assistance of the failing left ventricle (LV) can lead to functional recovery after a period of unloading, including restoration of β -adrenergic receptor (β AR) inotropic reserve. We tested whether prolonged LV unloading of failing rabbit hearts by use of a heterotopic transplantation technique could lead to recovery and whether adenoviral gene transfer of a β_2 AR transgene (Adv- β_2 AR) could alter this process.

Methods and Results—Heart failure was induced by coronary artery ligation in adult New Zealand White rabbits. After 4 weeks, failing hearts were heterotopically transplanted into recipient rabbits, allowing normal coronary perfusion but complete LV unloading. We also placed an LV latex balloon for remote access and in vivo physiological analysis. We found that there was reversal of signaling and functional abnormalities after 30 days of unloading. In another set of failing hearts, we randomly delivered, at the time of transplantation, either 2×10^{11} viral particles of Adv- β_2 AR or saline via the coronary arteries. Sham-operated animals with nonfailing hearts served as controls. After 5 days of unloading, in vivo LV contractility (LV dP/dt_{max}) and relaxation (LV dP/dt_{min}) were significantly decreased in saline-treated failing hearts compared with control nonfailing hearts ($P < 0.05$). In failing hearts treated with Adv- β_2 AR, however, LV dP/dt_{max} and LV dP/dt_{min} were improved in response to higher preloads ($P < 0.05$) and β AR stimulation ($P < 0.01$).

Conclusions—Heterotopic transplantation in the rabbit does allow recovery of the failing heart, and β_2 AR overexpression acutely enhances this functional improvement. Accordingly, genetic manipulation of β AR signaling may represent a novel molecular adjunct to mechanical assistance to facilitate functional myocardial recovery. (*Circulation*. 2002;106:124-129.)

Key Words: heart failure ■ receptors, adrenergic, beta ■ remodeling ■ heart-assist device ■ gene therapy

Alterations that take place in the β -adrenergic receptor (β AR) system during progression of heart failure (HF) are well characterized.¹ Typically, β_1 AR density is selectively reduced and the remaining β_1 - and β_2 ARs are desensitized.² Studies in transgenic mice have revealed that cardiac overexpression of β_2 ARs can be a mechanism for enhanced inotropic support for the heart.^{3,4} These studies have also been extended to normal rabbits in which intracoronary delivery of a β_2 AR-containing adenovirus (Adv- β_2 AR) has improved in vivo LV function,^{5,6} suggesting that overexpression of β_2 ARs may represent a novel strategy to improve ventricular function in patients with HF. In contrast to the apparent therapeutic effect of β_2 ARs, β_1 AR overexpression in the hearts of transgenic mice, even at low levels, leads to progressive early cardiomyopathy and HF.⁷

Treatment of patients with end-stage HF involves different modalities, including mechanical support with a left ventricular assist device (LVAD) and cardiac transplantation. Recently, significant improvement in ventricular function has been reported in some patients assisted with a mechanical

pump, and surprisingly, successful weaning from the LVAD was occasionally possible.⁸⁻¹² Little is known regarding the mechanisms that might predict this functional recovery,¹⁰ although contractile response to β AR stimulation improved in samples obtained from human ventricles unloaded with an LVAD.^{13,14} Therefore, a better understanding of the “reverse remodeling” process, ie, the mechanisms involved in such functional recovery during ventricular unloading, may lead to new molecular approaches that could provide novel hemodynamic support strategies to favor myocardial recovery and potentially successful weaning from LVADs without orthotopic transplantation.

In the present study, we wanted to determine whether heterotopic transplantation of failing rabbit hearts could represent a model of LV unloading and support functional recovery. Moreover, because β AR stimulation is the most powerful means of enhancing ventricular contractility, we investigated the potential benefit of combining genetic augmentation of β_2 AR signaling and ventricular unloading for the reverse remodeling process.

Received February 18, 2002; revision received April 17, 2002; accepted April 17, 2002.

From the Departments of Surgery (H.T.T., A.D.E., G.B.W., W.J.K.) and Medicine (K.W.), Duke University Medical Center, Durham, NC. Dr Tevæarai is now at the Department of Cardiovascular Surgery, Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland.

Correspondence to Walter J. Koch, PhD, Box 2606, Duke University Medical Center, Durham, NC 27710. E-mail koch0002@mc.duke.edu

© 2002 American Heart Association, Inc.

Circulation is available at <http://www.circulationaha.org>

DOI: 10.1161/01.CIR.0000020220.79105.FD

TABLE 1. Comparison of LV dP/dt_{max}, LV dP/dt_{min}, and LV Systolic Pressure in Failing (4 Weeks Post-MI) or Control (4 Weeks Post-Sham) Hearts at Day 1 After Transplantation

	HF (n=10)	Control Hearts (n=7)
LV dP/dt _{max} , mm Hg/s		
Baseline	707.5±83.8*	986.8±93.8
+0.3 mL	810.0±99.3†	1342.2±99.2
ISO+0.3 mL	1200.6±134.7†	2075.6±203.4
LV dP/dt _{min} , mm Hg/s		
Baseline	-718.6±67.7*	-945.7±102.9
+0.3 mL	-853.2±88.8†	-1302.3±74.4
ISO+0.3 mL	-1208.7±138.3†	-1916.1±151.6
LV systolic pressure, mm Hg		
Baseline	42.8±2.9	50.9±5.6
+0.3 mL	70.0±5.2*	86.3±7.6
ISO+0.3 mL	78.8±7.0*	108.8±10.8

Baseline represents conditions in which the LV balloon and volume was adjusted to an LV end-diastolic pressure of 0 mm Hg; +0.3 mL, a volume of 0.3 mL; and ISO+0.3 mL, both ISO (0.1 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) and a 0.3 mL increased preload.

* $P < 0.05$, † $P < 0.005$ vs control hearts.

Methods

Adenoviral Transgenes

We used a first-generation E1/E3-deleted replication-deficient serotype 5 adenovirus as previously described.^{5,15} The β_2 AR transgene (Adv- β_2 AR) and the marker transgene β -galactosidase (Adv- β Gal) were driven by the cytomegalovirus promoter. Virus was thawed and mixed with PBS to a final volume of 2 mL immediately before use.

Induction of HF

Adult male New Zealand White rabbits (3 kg) were used, and all procedures were performed in accordance with the regulations adopted by the National Institutes of Health and approved by the Animal Care and Use Committee of Duke University. A myocardial infarction (MI) was induced by a ligation of the first marginal branch of the left circumflex coronary artery as previously described.^{16,17} The control group included sham-operated animals in which only a thoracotomy and pericardiotomy were performed.

Heterotopic Transplantation and Gene Delivery

Four weeks after left circumflex coronary artery ligation, donor animals were anesthetized with ketamine (60 mg/kg) and acepromazine (1.0 mg/kg) and ventilated. The hearts were exposed via a clam-shell thoracic incision. HF was confirmed by global dilation of the heart, the presence of pleural effusion and/or ascites, and the presence of a large infarcted area estimated to be $\geq 30\%$ of the LV free wall. The donor hearts were then arrested by intracoronary perfusion with 30 mL of University of Wisconsin cold cardioplegic solution and quickly harvested and maintained at 4°C as described.^{18,19} Explanted hearts were randomized to receive either adenovirus or PBS. The hearts were then transplanted into the neck of recipient rabbits as described.¹⁸ Total ischemic time was ≈ 45 minutes. The hearts resumed vigorous contraction within 3 minutes of reperfusion. Dexamethasone (4 mg/kg) was administered intravenously before reperfusion and then intramuscularly on a daily basis. In a subgroup, an LV biopsy was taken from the donor heart for biochemical comparison with paired samples obtained 5 days after unloading.

LV Functional Assessment

In another subgroup of experiments, we positioned an LV latex balloon into the left atrium connected to tubing conducted under the skin to the subscapular region. To measure cardiac function, animals

were lightly anesthetized, and the extremity of the tubing was retrieved and connected to a Y-connector that allowed adjustment of the LV end-diastolic volume from one port and the introduction of a high-fidelity pressure transducer (Millar Instruments) through the other port. Baseline LV end-diastolic volume was normalized to a balloon volume providing an end-diastolic pressure of 0 mm Hg. LV function was assessed at days 1, 3, and 5 under 2 different preload conditions (baseline and +0.3 mL). Response to β AR stimulation was assessed by an intravenous infusion of 0.1 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ isoproterenol (ISO). At day 5, animals were euthanized, and infarction size was calculated as a percentage of the entire free wall as described.¹⁷ In a separate group of animals, longer-term effects of unloading were also examined by measurement of LV function every 5 days until day 30.

Determination of Myocardial β AR Density

Cardiac sarcolemmal membranes were prepared and total β AR density was determined as previously described.^{3,5}

Statistical Analysis

Data are expressed as mean \pm SEM. Paired Student's *t* tests were used for comparison of β AR density before and after transplantation.




Figure 1. Progression of LV recovery over a 30-day period of unloading of failing hearts (4 weeks after MI, n=5). A, LV contractility (LV dP/dt_{max}). B, LV relaxation (LV dP/dt_{min}). Basal conditions were defined as the LV end-diastolic volume necessary to generate an LV end-diastolic pressure of 0 mm Hg. Basal function (\diamond) did not improve over the 30-day unloading period (* $P = \text{NS}$; 1-way ANOVA for repeated measures). Response to a perfusion of 0.1 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ISO (\triangle) improved slightly over time ($\dagger P < 0.05$ for LV dP/dt_{max}, $P = \text{NS}$ for LV dP/dt_{min}). The function in loaded LV (basal+0.3 mL end-diastolic volume) improved significantly over time (\square ; $\ddagger P < 0.05$ for LV dP/dt_{max}, $P < 0.01$ for LV dP/dt_{min}). ISO stimulation (0.1 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) of loaded LV (basal+0.3 mL end-diastolic volume) improved to the greatest extent over the 30-day period of LV unloading (\blacksquare ; $\S P < 0.001$ for LV dP/dt_{max} and for LV dP/dt_{min}). Txp indicates transplantation.

One-way and 2-way ANOVAs were used to compare LV function in transplanted hearts. For all tests, a value of $P < 0.05$ was considered significant.

Results

Assessment of HF

A total of 42 rabbit hearts were successfully transplanted into the neck of recipient animals and studied regularly until euthanization at day 5 or day 30. Hearts with an infarct size estimated to be $< 30\%$ of the LV free wall (at the time of death) and that were not accompanied by ascites and/or pleural effusion were not considered for transplantation. Importantly, no significant difference in infarct size was observed between MI hearts that received Adv- β_2 AR ($35.0 \pm 2.3\%$ of the LV free wall, $n=4$) and MI hearts treated with PBS ($38.9 \pm 1.9\%$; $n=6$, $P=NS$).

In vivo LV hemodynamics was measured in 4-week post-MI hearts via the inserted latex balloon at day 1 after transplantation. This function 24 hours after transplant allowed the effect of cardioplegic arrest and surgery to become less important.^{19,20} All indices of LV function were significantly decreased compared with the function of sham-operated transplanted hearts ($n=7$), confirming that these hearts were in HF (Table 1).

Functional Recovery of Unloaded (Transplanted) Failing Hearts

We tested whether chronic unloading via heterotopic transplantation could lead to reversal of LV dysfunction. To aid against rejection,¹⁸ we treated rabbits with a daily dose of dexamethasone, and LV function was assessed every 5 days until day 30 after surgery. Basal LV function did not change over the 30-day period of unloading in hearts transplanted 4 weeks after MI. In post-MI hearts, however, both LV contractility (LV dP/dt_{max}) and relaxation (LV dP/dt_{min}) progressively increased after ISO stimulation during the 4 weeks of unloading (Figure 1). Moreover, the response to an increased preload also improved in these failing hearts during the 30 days of unloading, and interestingly, the combination of both

increased preload and ISO provided the highest improvement over the 30 days of ventricular unloading (Figure 1).

Restoration of β AR Density in Unloaded Failing Ventricles

As in other models of HF, β AR density in LV biopsies obtained at the time of transplantation of the failing hearts (post-MI) was decreased (31.8 ± 2.6 fmol/mg protein, $n=6$) compared with control (sham) (46.4 ± 2.2 fmol/mg protein, $n=4$, $P < 0.005$) (Figure 2). Interestingly, β AR density was restored to normal levels (48.1 ± 3.3 fmol/mg protein, $n=6$, $P < 0.0005$ versus pretransplantation) in membranes purified from the same failing hearts after 5 days of unloading via heterotopic transplantation (Figure 2).

Transgene Expression

Five days after ex vivo delivery of 2×10^{11} total viral particles (TVP) of Adv- β Gal, the transgene was robustly expressed in the transplanted heart, as confirmed by histological galactosidase staining (Figure 3A). Expression of Adv- β_2 AR was dose-dependent, and both 1×10^{11} ($1 \times$) and 2×10^{11} ($2 \times$) TVP significantly increased β AR density, with the $2 \times$ dose




Figure 2. Comparison of β AR density in membrane (mb) preparations obtained from LV before heterotopic transplantation (pre-Txp) and 5 days after LV unloading (post-Txp) in normal (□; $n=4$) and failing (■, $n=6$) hearts. In normal hearts, β AR density increased from 46.4 ± 2.2 to 55.8 ± 2.7 fmol/mg mb protein ($*P < 0.05$). β AR density was decreased significantly in failing hearts before transplantation (31.8 ± 2.6 fmol/mg mb protein; $\dagger P = 0.002$) but increased to 48.1 ± 3.3 fmol/mg mb protein after 5 days of unloading ($\ddagger P < 0.0001$), which was similar to normal unloaded hearts ($\$P = NS$).





Figure 3. Myocardial transgene expression 5 days after cardiac heterotopic transplantation (Txp) and intracoronary gene delivery. A, β -Gal staining in a representative normal LV 5 days after Txp and delivery of 2×10^{11} TVP of Adv- β Gal. B, β AR density in membrane (mb) preparation obtained from Txp failing hearts and comparison between 3 doses of Adv- β_2 AR: 0.3×10^{11} TVP ($n=6$), 1×10^{11} TVP ($n=3$), and 2×10^{11} TVP ($n=4$). Control hearts received PBS only ($n=6$). $*P < 0.001$ vs control.

TABLE 2. Comparison of LV dP/dt_{max}, LV dP/dt_{min}, and LV Systolic Pressure 5 Days After LV Unloading Between Failing Hearts (4 Weeks Post-MI) That Received Either PBS (HF+PBS) or Adv-β₂AR (HF+βAR) at the Time of Transplantation, and Control Hearts (4 Weeks Post-Sham) That Received PBS (Control Hearts+PBS)

	HF+PBS (n=6)		HF+βAR (n=4)		Control Hearts+PBS (n=7)	
	Day 1	Day 5	Day 1	Day 5	Day 1	Day 5
LV dP/dt_{max}						
Baseline	678.5±97.2	849.3±120.8	909.5±170.0	1349.1±377.8	986.8±93.8	1527.4±142.4†
+0.3 mL	806.6±122.7	935.0±139.0	1021.3±218.2	2224.8±449.0*	1342.2±99.2	2289.8±135.3‡
ISO+0.3 mL	1253.5±169.6	1350.1±175.5	1548.2±409.5	2652.6±393.9*	2075.6±203.4	3230.5±222.8‡
LV dP/dt_{min}						
Baseline	-652.3±74.4	-764.3±93.5	-942.8±109.9	-1157.0±233.7	-945.7±102.9	-1218.1±95.7†
+0.3 mL	-834.1±104.8	-892.3±126.9	-1026.1±170.4	-1623.7±271.0*	-1302.3±74.4	-1879.0±70.7‡
ISO+0.3 mL	-1257.4±180.5	-1203.7±153.2	-1459.6±362.2	-1965.5±229.2*	-1916.1±151.6	-2457.4±115.9‡
LV systolic pressure						
Baseline	41.5±3.8	45.7±3.6	53.1±7.3	68.4±15.1	50.9±5.6	76.1±6.8†
+0.3 mL	65.4±6.8	68.9±6.5	85.5±7.9	139.0±14.8†	86.3±7.6	149.5±9.6‡
ISO+0.3 mL	75.7±10.6	77.7±10.4	103.5±17.1	158.1±16.7†	108.8±10.8	163.7±8.5‡

Conditions and units are identical to Table 1.

* $P<0.05$, † $P<0.01$, ‡ $P<0.001$ vs group of HF+PBS.

increasing it to 511.6 ± 101.5 fmol/mg membrane protein (n=4). This represents >8 times the value measured in control hearts (Figure 3B; $P<0.005$). Accordingly, we used 2×10^{11} TVP as our unloaded heart dose.

β₂AR Gene Transfer Improves Functional Recovery of Unloaded Failing Hearts

Although baseline function did not differ significantly between failing hearts that received the β₂AR transgene (n=4) and those that received PBS (n=6), we found an improved response to mechanical stimulation when we increased preload to 0.3 mL above baseline volume ($P<0.05$, Table 2 and Figure 4). We also observed a significant functional improvement in response to βAR stimulation after 5 days of LV unloading ($P<0.05$, Table 2 and Figure 4). Moreover, maximal LV pressure generated during systole was significantly increased in unloaded failing hearts previously treated with Adv-β₂AR compared with those that received PBS ($P<0.005$, Figure 5). In fact, functional response to ISO stimulation was virtually identical to that obtained from nonfailing (sham, n=7) unloaded hearts (Figure 5B).

Restoration of βAR Density Is Independent of Cardiac Denervation

Because the transplantation technique renders the grafted heart denervated, restoration of βAR density might be a result of the absence of direct neural stimulation rather than unloading. To test this, we compared the function of normal transplanted hearts in which the LV balloon was kept inflated to maintain LV loading with that of unloaded transplanted normal hearts in which the balloon was deflated, except for data recordings at day 1 and day 5. Although no baseline difference was observed between loaded and unloaded hearts after 5 days, both βAR and preload-induced function progressively improved in unloaded hearts, whereas no change occurred in loaded hearts (Figure 6A). In addition, βAR

density significantly increased in unloaded hearts compared with nontransplanted hearts, whereas no difference was observed in transplanted hearts in which the LV balloon was kept inflated (Figure 6B). Thus, restoration of βAR density as well as functional improvement appears to be due to mechanical unloading rather than denervation.

Discussion

In the present study, we used the heterotopic transplantation model to simulate ventricular unloading. Using this model, we were able to reproduce the functional improvements observed in human failing hearts unloaded by use of an LVAD. Moreover, we demonstrate the benefit of gene-




Figure 4. Comparison of contractility between day 1 (□) and day 5 (■) after heterotopic transplantation of normal hearts (Ct; n=7), failing hearts treated by transplantation and treated with Adv-β₂AR (HF+β; n=4), and failing transplanted hearts that received PBS (HF; n=6). Shown is contractility at baseline (LV end-diastolic pressure=0 mm Hg), to increased preload conditions (+0.3 mL), and after βAR stimulation with $0.1 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ISO, and under a combination (ISO+0.3 mL). Results are expressed as percentage of the basal value measured at day 1 after transplantation. * $P=NS$, † $P<0.05$, ‡ $P<0.01$ (day 5 vs day 1).




Figure 5. LV systolic maximal pressure (5 days after transplantation) in response to increased preload conditions and (A) without or (B) with ISO stimulation ($0.1 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) in normal hearts (Δ ; $n=7$) and failing hearts that received Adv- β_2 AR (\blacksquare ; $n=4$) or PBS (\blacklozenge ; $n=6$). * $P<0.05$, † $P<0.005$ (Adv- β_2 AR vs PBS). ‡ $P=NS$ (ANOVA, Adv- β_2 AR transplanted hearts vs normal transplanted hearts).

mediated modulation of β_2 ARs to restore functional response to β AR stimulation as well as to increased preload. Thus, our data indicate that it may be advantageous to combine mechanical unloading with a molecular therapeutic modality to hasten and enhance myocardial functional recovery. As described previously, not all patients may recover sufficient ventricular strength to allow weaning from their LVAD^{8–12} and prolonged recovery of their myocardial contractile function.²¹ In the present study, failing hearts required 1 month of unloading after a large LV infarction before we were able to measure functional improvement. In contrast, failing hearts that received intracoronary Adv- β_2 AR demonstrated significant LV functional improvement within the first week of ventricular unloading. Thus, Adv- β_2 AR gene therapy to the unloaded failing heart may represent a novel form of “molecular assistance” to provide beneficial synergy to mechanical assistance.

Previous reports have suggested a role for β AR signaling in the reverse remodeling process that occurs in mechanically assisted failing hearts.^{13,14} For example, response to β AR stimulation was increased in human myocytes isolated from hearts unloaded with an LVAD.^{13,14} In the present study, β AR density was increased in normal hearts 5 days after heterotopic transplantation, and we verified that these changes were independent of denervation. Five days was not sufficient, however, to permit any significant functional improvement due to unloading. Importantly, overexpression of β_2 ARs to ≈ 8 -fold over normal via adenovirus-mediated




Figure 6. Contractility and β_2 AR density in chronically loaded and unloaded control transplanted hearts. A, LV dP/dt_{max} and LV dP/dt_{min} of normal hearts 5 days after transplantation. Comparison is made between unloaded (\square ; $n=5$) and loaded (\blacksquare ; $n=3$) hearts as described in the text. B, β_2 AR density in these loaded ($n=3$) vs unloaded ($n=3$) transplanted hearts. Values are expressed as percentage of a control group of normal nontransplanted hearts ($n=3$). * $P<0.005$ (unloaded hearts vs control nontransplanted hearts). † $P=NS$ (loaded hearts vs control nontransplanted hearts).

β_2 AR gene delivery at the time of transplantation was accompanied by significant increases in both contractility and relaxation in response to preload and β AR agonist stimulation within just 5 days. Thus, manipulation of β_2 AR signaling appears to be able to play a critical role in the reverse remodeling process by triggering functional recovery of unloaded failing hearts.

It is noteworthy that in transgenic mice, cardiac overexpression of β_2 ARs at extraordinary high levels (>150 -fold) was associated with cardiomyopathy, whereas cardiac β_2 AR overexpression to up to ≈ 100 -fold showed no pathology.^{3,4} These studies in transgenic

mice led to the hypothesis that β_2 AR overexpression would be beneficial in HF. Indeed, β_2 AR gene delivery to normal rabbit hearts also enhanced function, although β_2 AR overexpression was much lower.^{5,6} Moreover, β_2 AR overexpression, at levels seen in the present study, in cultured failing rabbit cardiomyocytes did improve abnormal β AR signaling; however, functional indices were not measured. Importantly, actual treatment of a failing heart with β_2 AR gene transfer, such as in the present study, has not previously been performed. In this study as well as in our previous adenovirus-mediated gene delivery studies, we have not seen cardiac β_2 AR overexpression exceed ≈ 15 -fold.^{5,6} Thus, we do not believe that this methodology would lead to “toxic” levels of β_2 AR overexpression, but importantly, we appear to be able to achieve “therapeutic” levels of β_2 ARs in the heart.

It is important to note the appreciation that signaling through β_1 ARs and β_2 ARs are qualitatively and quantitatively different, which was reviewed recently.²² In contrast to the beneficial cardiac effects after β_2 AR overexpression (up to 100-fold), transgenic overexpression of β_1 ARs in the mouse heart even at low levels (15- to 30-fold) produces significant pathology, including cardiomyopathy and HF. Moreover, some data indicate that these 2 β AR subtypes can exhibit opposing effects in the heart, including the induction of apoptosis.²² Thus, as this study indicates, selective enhancement of β_2 AR signaling may be therapeutic in the failing heart, because we have shown that β_2 AR overexpression can hasten the functional recovery of unloaded failing hearts.

This study does present some limitations. We did not specifically address the arrhythmic activity in the transplanted hearts that may be expected after β_2 AR overexpression. Another concern in the interpretation of these results is immune rejection, although no significant signs of immune rejection were detected after 5 days of transplantation. Inflammatory reactions might trigger adverse reactions that could also influence the interactions between reverse remodeling, adenovirus-mediated modification of genetic expression, and subsequent functional changes. In addition, the nature of adenoviruses is such that they allow gene expression for a limited period of time and therefore may not be appropriate for long-term studies of reverse remodeling. This could be an advantage, however, because permanent overexpression may be deleterious after adequate functional recovery is achieved. Finally, in this model of unloading, the failing heart is removed from its environment and placed into a healthy recipient. Although we demonstrated that functional and β AR density recoveries were independent of cardiac denervation, other humoral molecules present in the HF patient may influence ventricular remodeling.

Nevertheless, our study clearly demonstrates the potential usefulness of β_2 AR augmentation to aid in the reverse remodeling process that occurs in unloaded hearts, because this may represent a beneficial molecular adjunct to mechanical unloading to improve functional recovery of severely dysfunctional hearts. Moreover, our study demonstrates that heterotopic transplantation can reasonably assimilate mechanical unloading and may thus represent a suitable model to study the functional and biochemical effects of adjunct therapeutic modalities.

Acknowledgments

This work was supported by grants from the Swiss National Science Foundation 84NP-057501 and 32-65044.01 (Dr Tevearai), National Institutes of Health grants HL-59533 (Dr Koch) and HL-56205 (Dr Koch), and a medical student research fellowship from Howard Hughes Medical Institute (G.B. Walton).

References

1. Brodde OE. Beta-adrenoceptors in cardiac disease. *Pharmacol Ther.* 1993;60:405–430.
2. Bristow MR, Hershberger RE, Port JD, et al. β_1 and β_2 adrenergic receptor mediated adenylate cyclase stimulation in nonfailing and failing human ventricular myocardium. *Mol Pharmacol.* 1989;35:295–303.
3. Milano CA, Allen LF, Rockman HA, et al. Enhanced myocardial function in transgenic mice overexpressing the β_2 -adrenergic receptor. *Science.* 1994;264:582–586.
4. Liggett SB, Tepe NM, Lorenz JN, et al. Early and delayed consequences of β_2 -adrenergic receptor overexpression in mouse hearts: critical role for expression level. *Circulation.* 2000;101:1707–1714.
5. Maurice JP, Hata JA, Shah AS, et al. Enhancement of cardiac function after adenoviral-mediated in vivo intracoronary β_2 -adrenergic receptor gene delivery. *J Clin Invest.* 1999;104:21–29.
6. Shah AS, Lilly RE, Kypson AP, et al. Intracoronary adenovirus-mediated delivery and overexpression of the β_2 -adrenergic receptor in the heart: prospects for molecular ventricular assistance. *Circulation.* 2000;101:408–414.
7. Engelhardt S, Hein L, Wiesmann F, et al. Progressive hypertrophy and heart failure in β_1 -adrenergic receptor transgenic mice. *Proc Natl Acad Sci U S A.* 1999;96:7059–7064.
8. Frazier OH, Benedict CR, Radovancevic B, et al. Improved left ventricular function after chronic left ventricular unloading. *Ann Thorac Surg.* 1996;62:675–682.
9. Müller J, Wallakut G, Weng Y, et al. Weaning from mechanical support after complete recovery in patients with idiopathic dilated cardiomyopathy. In: Hetzer R, Hennig E, Loebe M, eds. *Mechanical Circulatory Support.* Berlin: Springer; 1997; 93:107.
10. Mancini DM, Beniaminovitz A, Levin H, et al. Low incidence of myocardial recovery after left ventricular assist device implantation in patients with chronic heart failure. *Circulation.* 1998;98:2383–2389.
11. Pietsch L, Laube H, Baumann G, et al. Recovery from end-stage ischemic cardiomyopathy during long-term LVAD support. *Ann Thorac Surg.* 1998;66:555–557.
12. Hetzer R, Müller J, Weng Y, et al. Cardiac recovery in dilated cardiomyopathy by unloading with a left ventricular assist device. *Ann Thorac Surg.* 1999;68:742–749.
13. Dipla K, Mattiello JA, Jeevanandam V, et al. Myocyte recovery after mechanical circulatory support in humans with end-stage heart failure. *Circulation.* 1998;97:2316–2322.
14. Heerd PM, Holmes JW, Cai B, et al. Chronic unloading by left ventricular assist device reverses contractile dysfunction and alters gene expression in end-stage heart failure. *Circulation.* 2000;102:2713–2719.
15. Akhter SA, Skaer CA, Kypson AP, et al. Restoration of β -adrenergic signaling in failing cardiac ventricular myocytes via adenoviral-mediated gene transfer. *Proc Natl Acad Sci U S A.* 1997;94:12100–12105.
16. Maurice JP, Shah AS, Kypson AP, et al. Molecular β -adrenergic signaling abnormalities in failing rabbit hearts after infarction. *Am J Physiol.* 1999;276:H1853–H1860.
17. White DC, Hata JA, Shah AS, et al. Preservation of myocardial β -adrenergic receptor signaling delays the development of heart failure following myocardial infarction. *Proc Natl Acad Sci U S A.* 2000;97:5428–5433.
18. Shah AS, White DC, Tai O, et al. Adenovirus-mediated genetic manipulation of the myocardial β -adrenergic signaling system in transplanted hearts. *J Thorac Cardiovasc Surg.* 2000;120:581–588.
19. Tevearai HT, Eckhart AD, Shotwell KF, et al. Ventricular dysfunction after cardioplegic arrest is improved after myocardial gene transfer of a β -adrenergic receptor kinase inhibitor. *Circulation.* 2001;104:2069–2074.
20. DiSesa VJ. Pharmacologic support for postoperative low cardiac output. *Semin Thorac Cardiovasc Surg.* 1991;3:13–23.
21. Hetzer R, Muller JH, Weng YG, et al. Midterm follow-up of patients who underwent removal of a left ventricular assist device after cardiac recovery from end-stage cardiomyopathy. *J Thorac Cardiovasc Surg.* 2000;120:843–853.
22. Rockman HA, Koch WJ, Lefkowitz RJ. Seven membrane spanning receptors and heart function. *Nature.* 2002;415:206–212.