

# Gene Transfer of Endothelial Nitric Oxide Synthase Improves Relaxation of Carotid Arteries From Diabetic Rabbits

D.D. Lund, PhD; F.M. Faraci, PhD; F.J. Miller, Jr, MD; D.D. Heistad, MD

**Background**—Diabetes mellitus is associated with impairment of NO-mediated vascular relaxation. The purpose of this study was to determine whether adenovirus-mediated gene transfer of endothelial NO synthase (eNOS) or Cu/Zn superoxide dismutase (SOD1) improves responsiveness to acetylcholine in alloxan-induced diabetic rabbits.

**Methods and Results**—After 8 weeks, plasma glucose was greater in diabetic rabbits ( $418 \pm 35$  mg/dL) (mean  $\pm$  SEM) than in normal rabbits ( $105 \pm 4$  mg/dL). Carotid arteries were removed and cut into ring segments. Arteries were incubated for 2 hours with adenoviral vectors driven by a CMV promoter expressing  $\beta$ -galactosidase ( $\beta$ -gal), eNOS, SOD1, or vehicle. After incubation with virus, arteries were incubated for an additional 24 hours to allow transgene expression. Vascular reactivity was examined by recording isometric tension. After precontraction with phenylephrine, responses to the endothelium-independent vasodilator sodium nitroprusside were similar in diabetic and normal arteries. Endothelium-dependent relaxation to acetylcholine ( $3 \times 10^{-6}$  mol/L) was significantly less in arteries from diabetic animals ( $68 \pm 5\%$ ) than in normal vessels ( $90 \pm 3\%$ ). Adenoviral transfection of arteries with eNOS improved relaxation in response to acetylcholine in diabetic ( $EC_{50}$  eNOS =  $0.64 \pm 0.12 \times 10^{-7}$  mol/L versus vehicle =  $1.70 \pm 0.43 \times 10^{-7}$  mol/L) but not normal arteries. Vasorelaxation in response to acetylcholine was inhibited by *N*<sup>o</sup>-nitro-L-arginine (100  $\mu$ mol/L) in all groups. Responses to acetylcholine were unchanged after gene transfection of SOD1 or  $\beta$ -gal in arteries from diabetic or normal rabbits.

**Conclusions**—Adenovirus-mediated gene transfer of eNOS, but not SOD, improves impaired NO-mediated relaxation in vessels from diabetic rabbits. (*Circulation*. 2000;101:1027-1033.)

**Key Words:** diabetes mellitus ■ acetylcholine ■ viruses ■ gene therapy ■ nitric oxide

Adenoviral vectors are a useful method for introducing genetic material into blood vessels to alter vascular function.<sup>1,2</sup> This approach has been used to improve function in blood vessels with functional impairment.<sup>3-7</sup>

Diabetes mellitus is often associated with impairment of endothelium-dependent relaxation in response to acetylcholine in humans<sup>8,9</sup> and experimental animals.<sup>10-14</sup> This observation suggests that an increase in vascular levels of endothelial NO synthase (eNOS), a major mediator of endothelium-dependent relaxation, might improve vascular dysfunction in diabetes. Thus, the first goal of this study was to determine whether gene transfer of eNOS improves vascular function in diabetic vessels.

Other studies indicate that endothelial dysfunction in diabetic vessels may be produced by excess production of reactive oxygen species.<sup>15,16</sup> Furthermore, administration of superoxide dismutase (SOD), which dismutates superoxide anion to H<sub>2</sub>O<sub>2</sub>, to diabetic animals improves endothelium-dependent relaxation.<sup>17,18</sup> This finding suggests that the dilator

response to acetylcholine might be improved by an increase in levels of SOD in vessels from diabetic animals. Thus, the second goal of this study was to determine whether gene transfer of SOD1 to carotid arteries from diabetic rabbits improves vascular function.

We have shown previously that after gene transfer, expression is greater in atherosclerotic than normal vessels<sup>19,20</sup> when a cytomegalovirus (CMV) promoter is used. Because the CMV promoter may be activated by reactive oxygen species, it seemed likely that expression of the transgene would be augmented in diabetic as well as atherosclerotic<sup>19,20</sup> vessels. The third goal of this study was to determine whether efficiency of gene transfer is augmented in diabetic vessels.

## Methods

### Adenovirus Vector

Three replication-deficient adenoviruses were used: (1) AdCMVeNOS (eNOS) constructed with cDNA for bovine eNOS,<sup>3</sup> (2) AdCMVSOD1 (SOD1) containing cDNA for human Cu/Zn-SOD,<sup>20</sup> and (3) AdCMV $\beta$ -galactosidase ( $\beta$ -gal) containing the reporter gene for  $\beta$ -gal, used as a

Received July 7, 1999; revision received August 25, 1999; accepted September 7, 1999.


From the Departments of Internal Medicine and Pharmacology and the Cardiovascular Center, University of Iowa College of Medicine and Veterans Affairs Medical Center, Iowa City.

Correspondence to Donald D. Heistad, MD, Department of Internal Medicine, University of Iowa College of Medicine, Iowa City, IA 52242. E-mail donald-heistad@uiowa.edu


© 2000 American Heart Association, Inc.

*Circulation* is available at <http://www.circulationaha.org>

A



B.



**Figure 1.** In situ detection of  $O_2^{\cdot -}$  in carotid artery. Confocal fluorescent photomicrographs of normal and diabetic rabbit carotid artery incubated with hydroethidine. Carotid artery from normal rabbit shows minimal fluorescence in endothelium (e) and adventitia. In contrast, carotid artery from diabetic rabbit shows increased ethidium bromide fluorescence reflecting increased  $O_2^{\cdot -}$  levels throughout vessel wall, which was greatest in endothelium. Similar findings were observed in 4 of 5 diabetic rabbits.

control virus.<sup>3,4</sup> A 3% sucrose PBS solution (vehicle) served as a virus-free control. Adenovirus was obtained from the Vector Core Laboratory of University of Iowa and stored at  $-80^\circ\text{C}$  until used.

### Animals

Hyperglycemia was produced in adult male New Zealand White rabbits (2.2 to 2.5 kg) by administration of alloxan (150 mg/kg IV) via the lateral ear vein. Rabbits that did not develop diabetes served as controls. At the end of 8 weeks, body weight increased in normal rabbits (n=6) from  $2.75\pm 0.06$  to  $3.73\pm 0.04$  kg and in diabetic rabbits (n=10) from  $2.68\pm 0.07$  to  $2.76\pm 0.15$  at the end of the study ( $P<0.05$  versus normal). Blood glucose concentrations at the end of the study were  $105\pm 4$  mg/dL in normal rabbits and  $418\pm 35$  mg/dL in diabetic animals ( $P<0.05$ , normal versus diabetic).

Eight weeks after induction of diabetes, rabbits were euthanized by injection of sodium pentobarbital (50 mg/kg) followed by exsanguination. The carotid arteries were quickly removed and placed in cold ( $4^\circ\text{C}$ ) oxygenated Krebs solution (mmol/L: NaCl 133, KCl 4.7,  $\text{NaH}_2\text{PO}_4$  1.35,  $\text{NaHCO}_3$  16.3,  $\text{MgSO}_4$  0.61, glucose 7.8, and  $\text{CaCl}_2$  2.52). The carotid arteries were then cut into segments 4 mm long for ex vivo incubation with adenovirus. All procedures and handling of animals were reviewed and approved by the Animal Care and Use Committee of the University of Iowa.

### Ex Vivo Infection

Rings from the carotid arteries were placed in a 96-well culture plate and incubated with either Ad5CMV $\beta$ -gal ( $1\times 10^9$  and  $3\times 10^9$  pfu/mL), Ad5CMV $\text{eNOS}$  ( $3\times 10^9$  pfu/mL), Ad5CMV $\text{SOD}$  ( $3\times 10^9$  pfu/mL), or vehicle (PBS with 3% sucrose) for 2 hours at  $37^\circ\text{C}$ . The rings were placed in Eagle's minimal essential media (Boehringer Mannheim) containing 100 U/mL penicillin and 100  $\mu\text{g}/\text{mL}$  streptomycin for an additional 24 hours at  $37^\circ\text{C}$  in a chamber aerated with 95%  $O_2$  and 5%  $\text{CO}_2$ .

### Detection of $\beta$ -Galactosidase


After ex vivo incubation, some vessels were rinsed with PBS and fixed with 2% paraformaldehyde and 0.2% glutaraldehyde in PBS for

detection of  $\beta$ -gal as previously described.<sup>3</sup> Vessel segments were incubated in 5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside (X-Gal, Sigma) solution for 2 hours at room temperature, embedded in paraffin, sectioned, and counterstained with nuclear fast red.


In other vessels, after incubation in adenovirus, the vessel segments were rinsed with PBS, frozen in liquid nitrogen, and stored at  $-70^\circ\text{C}$  until use.  $\beta$ -Gal activity was measured with a chemiluminescent reporter assay (Galacto-Light Plus, Tropix) as previously described.<sup>3</sup> Tissue was minced with a scalpel blade and placed in 150  $\mu\text{L}$  of Galacto-Light Lysis Solution (100 mmol/L potassium phosphate [pH 7.8], 0.2% Triton X-100). The homogenate was centrifuged at 10 000g for 10 minutes, and the supernatant was removed. The assay was performed with 10  $\mu\text{L}$  of supernatant in 200  $\mu\text{L}$  Galacton-Plus substrate:reaction buffer diluent (1:100 dilution). This reaction was carried out at room temperature, and light emissions were measured with a Moonlight 2010 luminometer (Analytical Luminescence Laboratory). A standard calibration curve was generated with purified *Escherichia coli*  $\beta$ -galactosidase (Boehringer Mannheim). Protein concentrations were determined with a Bio-Rad DC Protein Assay.  $\beta$ -Gal activity was expressed as mU  $\beta$ -gal/mg protein. Values for each group were calculated as an average of 2 rings from each animal.

### Detection of Superoxide

Hydroethidine, an oxidative fluorescent dye, was used to evaluate levels of superoxide in situ as described previously.<sup>21</sup> Cells are permeable to hydroethidine, and in the presence of  $O_2^{\cdot -}$ , hydroethidine is oxidized to fluorescent ethidium bromide, in which form it is trapped by intercalation with DNA. This method provides sensitive detection of  $O_2^{\cdot -}$  levels in situ. Unfixed frozen ring segments were cut into sections 30  $\mu\text{m}$  thick and placed on glass slides. Hydroethidine ( $2\times 10^{-6}$  mol/L) was applied to each tissue section and coverslipped. Slides were incubated in a light-protected humidified chamber at  $37^\circ\text{C}$  for 30 minutes. Images were obtained with a Bio-Rad MRC-1024 laser scanning confocal microscope equipped with a krypton/argon laser. Fluorescence was detected with a 585-nm long-pass filter. Normal and diabetic tissues were processed and



**Figure 2.** Histochemical staining of a carotid artery from a diabetic rabbit transfected with AdCMV  $\beta$ -gal. A, Carotid artery at  $\times 100$ ; B, carotid artery at  $\times 400$ . Blue color after X-gal staining represents expression of  $\beta$ -gal. Sections were counterstained with nuclear fast red.  $\beta$ -Gal is observed in endothelium and adventitia.



**Figure 3.** Effect of AdCMV $\beta$ -gal on  $\beta$ -gal activity (mU/mg protein) in carotid artery from control (open bars) and diabetic (solid bars) rabbits. Vessels were incubated with  $1 \times 10^9$  and  $3 \times 10^9$  pfu/mL of AdCMV $\beta$ gal. Values are mean  $\pm$  SEM, n=6.

imaged in parallel. Laser settings were identical for acquisition of images from normal and diabetic specimens.

Superoxide levels were also measured by lucigenin-enhanced chemiluminescence as described previously.<sup>21</sup> Vessel segments were placed in 0.5 mL PBS and lucigenin (0.25 mmol/L), and relative light units (RLUs) were measured with a Monolight 2010 luminometer. NADH oxidase has been proposed to be a major source of  $O_2^{\cdot -}$  in normal and diseased blood vessels.<sup>22</sup> To assess NADH oxidase activity in vessels, NADH (0.1 mmol/L) was added to the vessel segments, and  $O_2^{\cdot -}$  was measured by chemiluminescence for 5 minutes. Background counts were determined and subtracted, and RLUs were normalized to surface areas.

### Measurement of Vascular Function


Isometric tension was recorded to assess function of transfected carotid vessels. Vascular rings were mounted on stainless steel hooks at optimal resting tension (3 g) in individual organ baths bathed in Krebs bicarbonate solution at 37°C and aerated with 95%  $O_2$ /5%  $CO_2$ . Tension was periodically adjusted to the desired level during a 45-minute equilibration period. The vascular rings were then contracted twice with 100 mmol/L KCl and rinsed 3 times after each contraction. Responses to phenylephrine ( $10^{-9}$  to  $10^{-5}$  mol/L) were then examined. Concentration-response curves for the endothelium-dependent dilators acetylcholine ( $10^{-9}$  to  $10^{-5}$  mol/L) and A23187 ( $10^{-9}$  to  $10^{-5}$  mol/L) or the endothelium-independent dilator sodium nitroprusside ( $10^{-9}$  to  $10^{-5}$  mol/L) were then generated after precontraction of vessels with an  $EC_{50}$  dose of phenylephrine. In a separate set of vessels, responses to acetylcholine were examined in the presence of *N*<sup>o</sup>-nitro-L-arginine (L-NNA, 100  $\mu$ mol/L) to inhibit eNOS.

### Effects of Gene Transfer on Vascular Responses

	Phenylephrine		Acetylcholine		Nitroprusside		A23187	
	$ED_{50}, \times 10^{-6}$ mol/L	Peak Response, %	$ED_{50}, \times 10^{-7}$ mol/L	Peak Response, %	$ED_{50}, \times 10^{-8}$ mol/L	Peak Response, %	$ED_{50}, \times 10^{-8}$ mol/L	Peak Response, %
<b>Normal (n=6)</b>								
Vehicle	1.72 $\pm$ 0.22	117 $\pm$ 8	0.75 $\pm$ 0.10	90 $\pm$ 3	4.62 $\pm$ 0.79	98 $\pm$ 2	6.89 $\pm$ 1.10	91 $\pm$ 2
$\beta$ -gal	1.41 $\pm$ 0.25	119 $\pm$ 7	0.92 $\pm$ 0.23	84 $\pm$ 4	4.09 $\pm$ 0.45	97 $\pm$ 1	4.51 $\pm$ 0.52	87 $\pm$ 3
eNOS	1.39 $\pm$ 0.22	131 $\pm$ 9	0.93 $\pm$ 0.23	86 $\pm$ 2	3.34 $\pm$ 0.50	97 $\pm$ 1	4.78 $\pm$ 0.23	88 $\pm$ 2
SOD1	1.56 $\pm$ 0.29	127 $\pm$ 7	1.08 $\pm$ 0.19	87 $\pm$ 3	3.62 $\pm$ 0.38	98 $\pm$ 2	5.69 $\pm$ 0.63	89 $\pm$ 2
<b>Diabetic (n=10)</b>								
Vehicle	2.32 $\pm$ 0.51	151 $\pm$ 16	1.70 $\pm$ 0.43	68 $\pm$ 5 $\ddagger$	4.72 $\pm$ 1.24	96 $\pm$ 2	6.73 $\pm$ 1.60	85 $\pm$ 5
$\beta$ -gal	1.80 $\pm$ 0.39	144 $\pm$ 13	1.57 $\pm$ 0.23	62 $\pm$ 5 $\ddagger$	4.64 $\pm$ 0.74	95 $\pm$ 2	5.37 $\pm$ 0.81	76 $\pm$ 7
eNOS	2.19 $\pm$ 0.53	165 $\pm$ 20	0.64 $\pm$ 0.12* $\ddagger$	77 $\pm$ 4 $\ddagger$	4.05 $\pm$ 1.46	96 $\pm$ 2	6.89 $\pm$ 2.23	81 $\pm$ 5
SOD1	1.89 $\pm$ 0.29	153 $\pm$ 17	1.35 $\pm$ 0.31	71 $\pm$ 5 $\ddagger$	5.06 $\pm$ 1.40	95 $\pm$ 4	7.61 $\pm$ 1.80	80 $\pm$ 6

Data are mean  $\pm$  SEM.

\* $P=0.05$  vs vehicle-diabetic;  $\ddagger P=0.05$  vs  $\beta$ -gal-diabetic;  $\ddagger P=0.05$  vs corresponding group in normal rabbits.



**Figure 4.** Effect of gene transfer on response of carotid arteries to sodium nitroprusside in (A) normal rabbits (n=6) and (B) diabetic rabbits (n=10). Vessels were treated with  $\beta$ -gal or eNOS. Each point represents mean  $\pm$  SEM.

### Chemicals

Acetylcholine chloride, L-phenylephrine hydrochloride, sodium nitroprusside, L-NNA, and lucigenin were obtained from Sigma Chemical Co and dissolved in normal saline. Hydroethidine was obtained from Molecular Probes Inc, suspended in DMSO at a concentration of  $10^{-2}$  mol/L, and stored in aliquots at  $-80^\circ C$  until use.


### Statistical Analysis

Contractile responses were expressed as percent contraction of response to 100 mmol/L KCl, and relaxation was expressed as percent relaxation to contraction produced by an  $EC_{50}$  dose of phenylephrine. All data are expressed as mean  $\pm$  SEM. Intergroup comparisons were performed with an independent 1-way ANOVA to test for difference among treatment groups, followed by Bonferroni's corrected *t* test. Comparisons between diabetic and normal groups were made with Student's paired *t* test. Differences were considered to be significant at a value of  $P < 0.05$ .

## Results

### Superoxide Production

Vessels from diabetic rabbits had increased  $O_2^{\cdot -}$  levels measured by hydroethidine fluorescence compared with nor-



**Figure 5.** Effect of gene transfer on response of carotid arteries to acetylcholine without and with pretreatment with L-NNA in (A) normal rabbits ( $n=6$ ) and (B) diabetic rabbits ( $n=10$ ). Vessels were treated with  $\beta$ -gal or eNOS. Each point represents mean  $\pm$  SEM.

mal vessels (Figure 1). The increase in basal  $O_2^{\cdot -}$  levels was observed in endothelial cells, media, and adventitia.

With lucigenin chemiluminescence, there was no detectable basal production of  $O_2^{\cdot -}$  in segments of carotid artery. Because previous studies suggest that NADH/NADPH oxidase is a major source of  $O_2^{\cdot -}$  production in vessels, we assessed  $O_2^{\cdot -}$  production in vessels from normal or diabetic rabbits that were stimulated by NADH oxidase.<sup>21</sup> Superoxide production in response to NADH (0.1 mmol) was  $>2$ -fold greater in carotid arteries from diabetic rabbits than in normal carotids ( $277 \pm 54$  versus  $117 \pm 15$  RLU  $\cdot$  min<sup>-1</sup>  $\cdot$  mm<sup>-2</sup>,  $P < 0.05$ ). These data suggest increased propensity to generate  $O_2^{\cdot -}$  levels in carotid arteries from diabetic animals during treatment with NADH.

### Expression of $\beta$ -Gal

Twenty-four hours after incubation with AdCMV $\beta$ -gal, rings from the carotid artery were analyzed histochemically for transgene expression. Positive staining for  $\beta$ -gal was noted in adventitial and endothelial cells but not in vascular muscle (Figure 2). There was no staining in vehicle-treated vessels. The activity of  $\beta$ -gal was similar in normal and diabetic rabbits when carotid arteries were incubated with  $\beta$ -gal, which indicates similar expression of transgene product (Figure 3).

### Vasomotor Responses

In normal vessels after gene transfer of  $\beta$ -gal, SOD1, or eNOS, vasomotor responses to phenylephrine (Table), sodium nitroprusside (Figure 4A, Table), and A23187 (Table) were not different from those of vehicle-treated vessels. Relaxation to acetylcholine was also similar in vessels transfected with  $\beta$ -gal, eNOS, or SOD1 and in vehicle-treated vessels (Figure 5A). Relaxation to acetylcholine was inhibited after pretreatment with L-NNA (100  $\mu$ mol/L) in all vessels (Figure 5A).

In carotid arteries from diabetic rabbits, phenylephrine produced dose-dependent contraction, which was not altered by transfection with either eNOS or SOD1 compared with  $\beta$ -gal- or vehicle-treated animals (Table). Maximal contrac-

tion in vessels from diabetic rabbits ( $151 \pm 16\%$ ) tended to be greater than that in normal rabbits ( $117 \pm 8\%$ ), but it did not achieve statistical significance. In vehicle-treated vessels, responses to sodium nitroprusside, an endothelium-independent vasodilator, were similar in diabetic and normal rabbits. Transfection with  $\beta$ -gal, eNOS, or SOD1 did not alter the response to nitroprusside in vessels from diabetic rabbits (Figure 4A). In addition, vascular responses to the calcium ionophore A23187, an endothelium-dependent vasodilator, were similar in vehicle-treated rings in diabetic and normal rabbits. Transfection with  $\beta$ -gal, eNOS, or SOD1 did not alter the response to A23187 in vessels from diabetic rabbits (Table).

In vehicle-treated rings, maximal relaxation to acetylcholine was significantly less in diabetic ( $68 \pm 5\%$ ; Figure 5B) than in normal ( $90 \pm 3\%$ ; Figure 5A) rabbits ( $P < 0.05$ ). Transfection of  $\beta$ -gal or SOD1 did not alter responses to acetylcholine. Relaxation to acetylcholine was augmented in vessels from diabetic rabbits transfected with eNOS compared with incubation with vehicle or  $\beta$ -gal (Figure 5B). The EC<sub>50</sub> for eNOS was significantly different from that for vehicle or  $\beta$ -gal (Table) in diabetic rabbits. Relaxation to acetylcholine was inhibited in all vessels after pretreatment of rings with L-NNA (Figure 5B).

### Discussion

The major finding of this study is that adenovirus-mediated gene transfer of eNOS to carotid arteries improves impaired NO-mediated responses to acetylcholine in diabetic rabbits. This is the first demonstration, to the best of our knowledge, of the use of gene transfer to improve impaired vascular function in vessels during diabetes.

Endothelium-dependent relaxation is impaired in humans with both type I, insulin-dependent diabetes mellitus<sup>8</sup> and type II, non-insulin-dependent diabetes mellitus.<sup>9</sup> Endothelium-dependent relaxation is also a hallmark of impaired vascular responses in genetic models of diabetes<sup>13,14</sup> or in animals in which diabetes is induced with either alloxan or streptozotocin.<sup>10–12,15–18</sup> In this study, we observed less relaxation of carotid artery in response to acetylcholine in diabetic than in normal rabbits. Relaxation to sodium nitroprusside, an endothelium-independent vasodilator, was similar in normal and diabetic animals, which suggests that impaired relaxation to acetylcholine in diabetic rabbits is not due to dysfunction of vascular smooth muscle. These studies instead suggest impaired endothelium-dependent relaxation.

There was a tendency (not statistically significant) for greater maximal contraction to KCl in vessels from diabetic animals, which might make it difficult to compare relaxation in these vessels. Sodium nitroprusside was given to compare with responses to acetylcholine and to address differences in baseline. The key finding in this study is that gene transfer of eNOS improves relaxation to acetylcholine but not to sodium nitroprusside. These results cannot be explained by a tendency for augmented contraction to KCl in vessels from diabetic animals.

We did not measure vascular reactivity in fresh vessels to determine whether there is a change when vessels are maintained in culture, but we have previously reported that

vascular function in normal arteries is not impaired by incubation in tissue culture for 24 hours.<sup>4</sup> It was important for us to demonstrate, however, that vascular reactivity is still abnormal in arteries from diabetic animals<sup>10–18</sup> after maintenance in tissue culture for 24 hours. We found that impairment of responses in arteries from diabetic animals was comparable to that described previously.<sup>10–18</sup>

We and others have been successful in transferring eNOS cDNA to blood vessels. Although these studies have shown alteration in function after overexpression of eNOS in normal<sup>4,23</sup> and diseased<sup>3,5,7</sup> blood vessels, the functional effects of gene transfer of eNOS to vessels from diabetic animals have not been examined. It seemed important to determine whether overexpression of eNOS in arteries from diabetic rabbits might produce functional changes.

In this study, mechanisms responsible for endothelial dysfunction during diabetes appeared to be specific for a receptor-dependent stimulus of NO release, because the response to the receptor-independent endothelium-dependent vasomotor relaxing agent calcium ionophore A23187 was not impaired. Other investigators have also noted that vascular responses to A23187 may not be impaired in diabetic animals.<sup>24,25</sup>

We observed that cells were stained for  $\beta$ -galactosidase in the endothelium and adventitia of vessels from normal and diabetic animals. Fibroblasts in culture have muscarinic receptors,<sup>26,27</sup> and it is possible that adventitial fibroblasts in vivo also contain muscarinic receptors and thus could release NO in response to acetylcholine after transduction with eNOS. Other studies indicate that recombinant eNOS in adventitial fibroblasts in the dog basilar artery can be activated by bradykinin.<sup>28</sup> We have observed, however, that in the carotid artery of rabbits, after transduction of adventitia with eNOS and after denudation of endothelium, A23187 but not acetylcholine produces vascular relaxation.<sup>4</sup> Thus, it appears that endothelium is necessary to produce relaxation in response to acetylcholine, even after gene transfer of eNOS, in the rabbit carotid artery.

Numerous studies suggest that the mechanism of impaired endothelium-dependent relaxation in diabetes and atherosclerosis may involve inactivation of NO by oxygen-derived free radicals.<sup>12,13,17–20,24,29</sup> Production of superoxide anion inactivates NO,<sup>30,31</sup> and dismutation of free radicals has generally<sup>17,18</sup> but not always<sup>32</sup> been shown to improve impaired endothelium-dependent relaxation in experimental models of diabetes.<sup>17–18</sup>

Some investigators have suggested that an increase in activity of cyclooxygenase and release of prostanoids may be important in vascular dysfunction in diabetes,<sup>33</sup> but other studies indicate that indomethacin fails to alter endothelium-dependent relaxation in diabetes.<sup>13,34</sup> We chose not to use indomethacin in this study and thus cannot rule out the possibility that cyclooxygenase and vasoconstrictor prostanoids contribute to endothelial dysfunction in arteries from diabetic rabbits. In this preparation, however, acetylcholine-induced relaxation in arteries from normal and diabetic rabbits was blocked by L-NNA, which indicates that relaxation is mediated by NO. More importantly, overexpression of eNOS in arteries from diabetic rabbits improved vascular

function, even though the mechanism(s) of vascular dysfunction is not entirely clear.

We found an increase in ethidium bromide fluorescence in the vessels from diabetic rabbits. Ethidium bromide appears to be specific for superoxide in the vessel, because neither hydroxyl radical, NO, peroxyxynitrite, H<sub>2</sub>O<sub>2</sub>, hypochlorite, nor singlet O<sub>2</sub> significantly oxidizes hydroethidine.<sup>35</sup> Polyethylene-glycolated SOD abolished ethidium bromide fluorescence in blood vessels, which confirms the specificity of the fluorescent signal for superoxide anion.<sup>21</sup> Although we found increased O<sub>2</sub><sup>-</sup> levels in vessels from diabetic animals, gene transfer of SOD1 failed to improve vasomotor response to acetylcholine. The finding may result from several factors. First, in other studies in which exogenous SOD improves endothelium-dependent relaxation, the enzyme has access throughout the vessel wall. Adenovirus-mediated gene transfer increases SOD in the endothelium and adventitia but not the media.<sup>21</sup> Thus, enhanced production of free radicals in the media, which we demonstrated with hydroethidine (Figure 1), may not be corrected by gene transfer to endothelium and adventitia. Second, SOD1, which is present in cytosol, may not be able to protect NO from O<sub>2</sub><sup>-</sup> if the reaction of these radicals occurs in the extracellular space. Third, it is possible that abnormal relaxation to acetylcholine may not involve reaction with O<sub>2</sub><sup>-</sup>. Our findings and those of others<sup>24,25</sup> that responses to A23187 are not impaired in vessels from the diabetic animals suggest that there may be a selective effect on receptor-mediated endothelium-dependent relaxation, which may not be mediated by reactive oxygen species.

In previous studies, we observed greater expression after gene transfer to arteries in atherosclerotic than normal rabbits when an adenovirus with a CMV promoter was used.<sup>19,20</sup> In the present study, we observed no difference in activity of  $\beta$ -galactosidase in carotid arteries from normal and diabetic rabbits. Although both atherosclerosis and diabetes may have elevated concentrations of oxygen-derived free radicals and thus would be expected to increase expression when the CMV promoter is used, we speculate that this mechanism may not be as active in diabetic vessels as in atherosclerotic vessels and thus might fail to enhance gene expression.

The goal of this study was to use gene transfer as a tool for vascular biology. The major reason for studying the effects of gene transfer *ex vivo* instead of *in vivo* is that multiple mechanisms can be studied in the same rabbits; eg, in this study, we compared vehicle with gene transfer of eNOS, CuZn-SOD, and 2 concentrations of  $\beta$ -gal in the carotid arteries of each rabbit. Using *in vivo* techniques, we would need to use vehicle in 1 carotid artery versus an intervention in the other carotid artery. Because the design is less efficient, many more rabbits must be studied. With a similar design, to accomplish the same goals, we would need  $\approx 3$  times as many rabbits, and the approach would be less sensitive because comparison would be between multiple animals. Thus, this *ex vivo* gene transfer approach seems appropriate for mechanistic studies. In addition, the absence of an immune response *in vitro* also makes this approach attractive for mechanistic studies of vascular biology. Nevertheless, because of the immune response to adenovirus, it will be necessary to study gene transfer to vessels *in vivo*, especially if the approach were to move toward gene therapy.

In summary, this study demonstrates that adenoviral gene transfer of eNOS can improve impaired vascular function in diabetic vessels. Although several mechanisms may contribute to this response to eNOS, gene transfer provides a novel approach to study diabetic arteries. Gene transfer of SOD failed to improve vascular function. The finding of increased production of superoxide anion in the media of the artery, to which gene transfer of SOD does not have access, suggests that increased generation of superoxide throughout the arterial wall may play an important role in vascular dysfunction associated with diabetes mellitus.

### Acknowledgments

This work was supported by NIH grants HL-16066, NS-24621, HL-14388, and HL-38901; funds from the Department of Veterans Affairs; and funds from the Carver Trust of the University of Iowa. We would like to thank Dr Beverly L. Davidson, Richard D. Anderson, and the University of Iowa Gene Transfer Vector Core for preparation of virus. We thank Pam Tompkins and Margaret Donohue for their technical assistance and Arlinda LaRose for secretarial assistance.

### References

- Nabel EG. Gene therapy for cardiovascular disease. *Circulation*. 1995; 91:541–548.
- Heistad DD, Faraci FM. Gene therapy for cerebral vascular disease. *Stroke*. 1996;27:1688–1693.
- Ooboshi H, Toyoda K, Faraci FM, Lang MG, Heistad DD. Improvement of relaxation in an atherosclerotic artery by gene transfer of endothelial nitric oxide synthase. *Arterioscler Thromb Vasc Biol*. 1998;18:1752–1758.
- Ooboshi H, Chu Y, Rios CD, Faraci FM, Davidson BL, Heistad DD. Altered vascular function following adenovirus-mediated overexpression of endothelial nitric oxide synthase. *Am J Physiol*. 1997;273:H265–H270.
- Onoue H, Tsutsui M, Smith L, Stelter A, O'Brien T, Katusic ZS. Expression and function of recombinant endothelial nitric oxide synthase gene in canine basilar artery after experimental subarachnoid hemorrhage. *Stroke*. 1998;29:1959–1966.
- Channon KM, Qian H, Neplioueva V, Blazing MA, Olmez E, Shetty GA, Youngblood SA, Pawloski J, McMohon T, Stamler JS, George SE. In vivo gene transfer of nitric oxide synthase enhances vasomotor function in carotid arteries from normal and cholesterol-fed rabbits. *Circulation*. 1998;98:1905–1911.
- Von de Leyen HE, Gibbons GH, Morishita R, Lewis NP, Zhang L, Nakajima M, Kaneda Y, Cooke JP, Dzau VJ. Gene therapy inhibiting neointimal vascular lesion: in vivo transfer of endothelial cell nitric oxide synthase gene. *Proc Natl Acad Sci U S A*. 1995;92:1137–1141.
- McNally PG, Watt PAC, Rimmer T, Burden AC, Hearnshaw JR, Thurston H. Impaired contraction and endothelium-dependent relaxation in isolated resistance vessels from patients with insulin-dependent diabetes mellitus. *Clin Sci*. 1994;87:31–36.
- McVeigh GE, Brennan GM, Johnston GD, McDermott BJ, McGrath LT, Henry WR, Andrews JW, Hayes JR. Impaired endothelium-dependent and independent vasodilatation in patients with type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia*. 1992;35:771–776.
- Cohen RA. Dysfunction of vascular endothelium in diabetes mellitus. *Circulation*. 1993;87(suppl V):V67–V76.
- Pieper GM, Gross GJ. Oxygen free radicals abolish endothelium-dependent relaxation in diabetic rat aorta. *Am J Physiol*. 1988;255:H825–H833.
- Kamata K, Miyata N, Kasuya Y. Impairment of endothelium-dependent relaxation and changes in levels of cyclic GMP in aorta from streptozotocin-induced diabetic rats. *Br J Pharmacol*. 1989;97:614–618.
- Meraji S, Joakody L, Senaratene MP, Thomson ABR, Kappagoda T. Endothelium-dependent relaxation in aorta of BB rat. *Diabetes*. 1987;36:978–981.
- Pieper GM, Siebeneich W, Moore-Hilton G, Roza AM. Reversal by L-arginine of a dysfunctional arginine/nitric oxide pathway in endothelium of the genetic diabetic BB rat. *Diabetologia*. 1997;40:910–915.
- Diederich D, Scopec J, Diederich A, Dai FX. Endothelial dysfunction in mesenteric resistance arteries of diabetic rats: role of free radicals. *Am J Physiol*. 1994;266:H1153–H1161.
- Chang KC, Chung SY, Chong WS, Suh JS, Kim SH, Noh HK, Seong BW, Ko HJ, Chun KW. Possible superoxide radical-induced alteration of vascular reactivity in aortas from streptozotocin-treated rats. *J Pharmacol Exp Ther*. 1993;266:992–1000.
- Pieper GM, Mei DA, Langenstroer P, O'Rourke. Bioassay of endothelium-derived relaxing factor in diabetic rat aorta. *Am J Physiol*. 1988;263:H676–H680.
- Hattori Y, Kawasaki H, Abe K, Kanno M. Superoxide dismutase recovers altered endothelium-dependent relaxation in diabetic rat aorta. *Am J Physiol*. 1991;261:H1086–H1094.
- Lund DD, Faraci FM, Ooboshi H, Davidson BL, Heistad DD. Adenovirus-mediated gene transfer is augmented in basilar and carotid arteries of heritable hyperlipidemic rabbits. *Stroke*. 1998;29:120–125.
- Ooboshi H, Rios CD, Chu Y, Christensen SD, Faraci FM, Davidson BL, Heistad DD. Augmented adenovirus-mediated gene transfer in atherosclerotic vessels. *Arterioscler Thromb Vasc Biol*. 1997;17:1786–1792.
- Miller FJ, Gutterman DD, Rios CD, Heistad DD, Davidson BL. Superoxide production in vascular smooth muscle contributes to oxidative stress and impaired relaxation in atherosclerosis. *Circ Res*. 1998; 82:1298–1305.
- Rajogopalan S, Kurz S, Munzel T, Tarpey M, Freeman BA, Griending KK, Harrison DG. Angiotensin II-mediated hypertension in the rat increases vascular superoxide production via membrane NADH/NADPH oxidase activation. *J Clin Invest*. 1996;97:1916–1923.
- Chen AFY, O'Brien T, Tsutsui M, Kinoshita H, Pompili VJ, Crotty TB, Spector DJ, Katusic ZS. Expression and function of recombinant endothelial nitric oxide synthase in canine basilar artery. *Circ Res*. 1997;80:327–335.
- Pieper GM, Siebeneich W, Rosa AM, Jordan M, Adams MB. Chronic treatment in vivo with dimethylthiourea, a hydroxyl radical scavenger, prevents diabetes-induced endothelial dysfunction. *J Cardiovasc Pharmacol*. 1996;28:741–745.
- Pieper G. Divergent actions of chronic insulin treatment in vivo versus acute treatment ex vivo on diabetic-induced endothelial dysfunction. *Life Sci*. 1997;60:371–376.
- Haddad EB, Rousell J, Mak JC, Barns PJ. Long-term carbachol treatment-induced down-regulation of muscarinic M2-receptor but not m2 receptor mRNA in human lung cell line. *Br J Pharmacol*. 1995;116:2027–2032.
- Koenig JA, Edwardson JM. Intracellular trafficking of muscarinic acetylcholine receptor: importance of subtype and cell type. *Mol Pharmacol*. 1996;49:351–359.
- Tsutsui M, Chen AF, O'Brien T, Crotty TB, Katusic Z. Adventitial expression of recombinant eNOS gene restores NO production in arteries without endothelium. *Arterioscler Thromb Vasc Biol*. 1998;18:1231–1241.
- Pagano PJ, Griswold MC, Ravel D, Cohen RA. Vascular action of the hypoglycaemic agent gliclazide in diabetic rabbits. *Diabetologia*. 1998; 41:9–15.
- Marshall J, Wei EP, Kontos HA. Independent blockade of cerebral vasodilation from acetylcholine and nitric oxide. *Am J Physiol*. 1988;255:H847–H854.
- Rubanyi GM, Vanhoutte PM. Oxygen-derived free radicals, endothelium and responsiveness of vascular smooth muscle. *Am J Physiol*. 1986;250: H815–H821.
- Heygate KM, Lawrence IG, Bennett MA, Thurston H. Impaired endothelium-dependent relaxation in isolated resistance arteries of spontaneously diabetic rats. *Br J Pharmacol*. 1995;116:3251–3259.
- Tesfamariam B, Jakubowski JA, Cohen RA. Contraction of diabetic rabbit aorta caused by endothelium-derived PGH<sub>2</sub>-TxA<sub>2</sub>. *Am J Physiol*. 1989;257:H1327–H1333.
- Pieper GM. Enhanced, unaltered and impaired nitric oxide mediated endothelium-dependent relaxation in experimental diabetes mellitus: importance of disease duration. *Diabetologia*. 1999;42:204–213.
- Bindokas VP, Jordan J, Lee CC, Miller RJ. Superoxide production in rat hippocampal neurons: selective imaging with hydroethidine. *J Neurosci*. 1996;156:1324–1336.

## Gene Transfer of Endothelial Nitric Oxide Synthase Improves Relaxation of Carotid Arteries From Diabetic Rabbits

D. D. Lund, F. M. Faraci, F. J. Miller, Jr and D. D. Heistad

*Circulation*. 2000;101:1027-1033

doi: 10.1161/01.CIR.101.9.1027

*Circulation* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2000 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circ.ahajournals.org/content/101/9/1027>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation* is online at:  
<http://circ.ahajournals.org/subscriptions/>