

# Aerobic Fitness Is Associated With Cardiomyocyte Contractile Capacity and Endothelial Function in Exercise Training and Detraining

Ole Johan Kemi, MSc\*; Per Magnus Haram, BSc\*; Ulrik Wisløff, PhD; Øyvind Ellingsen, MD, PhD

**Background**—Physical fitness and level of regular exercise are closely related to cardiovascular health. A regimen of regular intensity-controlled treadmill exercise was implemented and withdrawn to identify cellular mechanisms associated with exercise capacity and maximal oxygen uptake ( $\dot{V}O_{2\max}$ ).

**Methods and Results**—Time-dependent associations between cardiomyocyte dimensions, contractile capacity, and  $\dot{V}O_{2\max}$  were assessed in adult rats after high-level intensity-controlled treadmill running for 2, 4, 8, and 13 weeks and detraining for 2 and 4 weeks. With training, cardiomyocyte length, relaxation, shortening,  $Ca^{2+}$  decay, and estimated cell volume correlated with increased  $\dot{V}O_{2\max}$  ( $r=0.92, -0.92, 0.88, -0.84, 0.73; P<0.01$ ). Multiple regression analysis identified cell length, relaxation, and  $Ca^{2+}$  decay as the main explanatory variables for  $\dot{V}O_{2\max}$  ( $R^2=0.87, P<0.02$ ). When training stopped, exercise-gained  $\dot{V}O_{2\max}$  decreased 50% within 2 weeks and stabilized at 5% above sedentary controls after 4 weeks. Cardiomyocyte size regressed in parallel with  $\dot{V}O_{2\max}$  and remained (9%) above sedentary after 4 weeks, whereas cardiomyocyte shortening, contraction/relaxation- and  $Ca^{2+}$ -transient time courses, and endothelium-dependent vasorelaxation regressed completely within 2 to 4 weeks of detraining. Cardiomyocyte length, estimated cell volume, width, shortening, and  $Ca^{2+}$  decay and endothelium-dependent arterial relaxation all correlated with  $\dot{V}O_{2\max}$  ( $r=0.85, 0.84, 0.75, 0.63, -0.54, -0.37; P<0.01$ ). Multiple regression identified cardiomyocyte length and vasorelaxation as the main determinants for regressed  $\dot{V}O_{2\max}$  during detraining ( $R^2=0.76, P=0.02$ ).

**Conclusions**—Cardiovascular adaptation to regular exercise is highly dynamic. On detraining, most of the exercise-gained aerobic fitness acquired over 2 to 3 months is lost within 2 to 4 weeks. The close association between cardiomyocyte dimensions, contractile capacity, arterial relaxation, and aerobic fitness suggests cellular mechanisms underlying these changes. (*Circulation*. 2004;109:2897-2904.)

**Key Words:** exercise ■ myocytes ■ hypertrophy ■ contractility ■ endothelium

Several lines of evidence suggest that regular physical exercise not only improves fitness and aerobic capacity but also reduces morbidity and mortality. Maximal oxygen uptake ( $\dot{V}O_{2\max}$ ) has emerged as an important clinical reference after epidemiological studies identified it as a major independent predictor of cardiovascular morbidity and mortality.<sup>1,2</sup> For successful implementation into standard prevention and therapy, cellular and molecular mechanisms of positive health effects need to be identified.

The present study is based on the notion that  $\dot{V}O_{2\max}$  is closely related to myocardial function. Accordingly, changes in cardiomyocyte size and function parallel those observed in  $\dot{V}O_{2\max}$  when an exercise regimen is implemented and withdrawn. Whereas regular exercise is known to increase cardiomyocyte function and dimensions<sup>3-6</sup> as well as endothelium-dependent arterial relaxation,<sup>6-10</sup> the

response to detraining has yet to be defined. In humans, detraining decreases  $\dot{V}O_{2\max}$ , but myocardial and arterial effects remain unclear.<sup>11-13</sup> Reported regression of training-induced hypertrophy ranges from none<sup>13</sup> to 20% over a period of 3 weeks.<sup>14</sup> Experimental data suggest hypertrophy regression at varying time courses,<sup>3,15-17</sup> with no account on cardiomyocyte or arterial function. Thus, the aim of the present study was to assess the cardiomyocyte contractile capacity and endothelium-dependent arterial relaxation changes in animals undergoing a controlled program of exercise training and detraining. We report  $\dot{V}O_{2\max}$  and associated changes in cardiomyocyte contractile capacity and endothelial function during detraining after 10 weeks of regular exercise and analyze correlations derived from previous 2- to 13-week training experiments in our laboratory.<sup>6,18</sup>

Received December 11, 2003; revision received March 9, 2004; accepted March 9, 2004.

From the Department of Circulation and Medical Imaging, Norwegian University of Science and Technology, Trondheim, Norway (O.J.K., P.M.H., U.W., Ø.E.), and the Department of Cardiology, St Olav's Hospital, Trondheim, Norway (U.W., Ø.E.).


\*The first 2 authors contributed equally to this work.

Correspondence to Øyvind Ellingsen, Department of Circulation and Medical Imaging, Medical Technology Research Center, Olav Kyrres gate 3, N-7489 Trondheim, Norway. E-mail oyvind.ellingsen@medisin.ntnu.no

© 2004 American Heart Association, Inc.

*Circulation* is available at <http://www.circulationaha.org>

DOI: 10.1161/01.CIR.0000129308.04757.72



**Figure 1.** Time course of  $\dot{V}O_{2\max}$  during 10 weeks of training (TR10, DETR2, DETR4) and detraining for 2 weeks (DETR2,  $\approx 50\%$  decrease in  $\dot{V}O_{2\max}$ ) and 4 weeks (DETR4,  $\dot{V}O_{2\max}$  stabilized) with corresponding sedentary controls (SED10, SED2, SED4). Time course of training was similar for all groups. Data are mean  $\pm$  SD. TR/DETR vs sedentary: \* $P < 0.01$ , † $P < 0.05$ . End point vs baseline: ‡ $P < 0.01$ , § $P < 0.05$ .

## Methods

### Study Design

For training-detraining experiments, a total of 36 female adult Sprague-Dawley rats (Møllegaards Breeding Center Ltd, Skensved, Denmark) were included and maintained as previously described.<sup>5,6</sup> Rats were randomized into 6 groups of either training/detraining or sedentary control, with 6 rats in each group. Groups TR10, DETR2, and DETR4 performed training until  $\dot{V}O_{2\max}$  was stable at a high level for 3 consecutive weeks (which occurred after 10 weeks), and then remained sedentary (ie, detraining) for 2 or 4 weeks, respectively. TR10 rats were euthanized 24 hours after the last exercise bout. DETR2 rats were euthanized when  $\approx 50\%$  of the exercise-gained  $\dot{V}O_{2\max}$  was lost, which occurred after 2 weeks, and DETR4 rats after 4 weeks of detraining, when  $\dot{V}O_{2\max}$  had been stable at a low level for 3 consecutive weeks. Corresponding sedentary control groups were SED10, SED2, and SED4. Detrained animals were euthanized 1 week after the latest  $\dot{V}O_{2\max}$  test. Thus, during detraining, animals were tested once a week, and when DETR4 rats showed an  $\approx 50\%$  decrease, DETR2 were not tested but euthanized. DETR4 were euthanized 1 week after no further decrease occurred. In DETR2, 2 rats were removed because of poor running, together with 2 corresponding controls. The relationships between  $\dot{V}O_{2\max}$  and cellular properties during training were investigated with unexplored data from previous studies<sup>6,18</sup> (the data appear in Figure 2). The Norwegian Council for Animal Research approved the experimental protocols.

### $\dot{V}O_{2\max}$ and Training

$\dot{V}O_{2\max}$  was measured during treadmill running in a metabolic chamber as previously described<sup>5,6,18</sup> at the start of every week in training/detraining animals to adjust training intensity and before and after the training period in the sedentary group. Training rats performed interval running 1 h/d, 5 d/wk, on a 25° inclined treadmill. After a 20-minute warm-up at 50% to 60% of  $\dot{V}O_{2\max}$ , exercise intervals alternated between 8 minutes at 85% to 90% of  $\dot{V}O_{2\max}$  and 2 minutes at 50% to 60%. Sedentary rats performed treadmill running for 15 minutes on a flat treadmill at 0.15 m/s for 2 d/wk to maintain running skills, which did not yield any training response; previous experiments indicate that this intensity corresponds to  $\approx 45\%$  of  $\dot{V}O_{2\max}$ .

### Cardiomyocyte Contractility, Calcium Handling, and Dimensions

Left ventricular myocytes were isolated as previously described with a modified Krebs-Henseleit  $Ca^{2+}$ -free buffer.<sup>5</sup> Collagenase II (250

IU/mL, Worthington), BSA (Sigma Chemical), and  $CaCl_2$  stepwise to 1.2 mmol/L were subsequently introduced. Ventricles were weighed after perfusion. Cells attached to laminin-coated coverslips rested 1 hour in HEPES buffer before 20 minutes of loading with 2  $\mu$ mol/L fura 2-AM (Molecular Probes) and were placed into a cell chamber (37°C) on an inverted microscope (Diaphot-TMD, Nikon) and stimulated electrically as previously described.<sup>6,18</sup> A 500-Hz rotating mirror alternated excitation wavelength between 340 and 380 nm, and 510-nm fluorescence emission was counted with a photomultiplier (D-104, Photon Technology International) and expressed as the ratio of the 2 excitation wavelengths. Cell shortening and relaxation were analyzed with video/edge-detection (model 104, Crescent Electronics). Ten stable, consecutive contractions at each stimulation frequency (2, 5, 7, and 10 Hz, and thereafter at 1 Hz to ensure that cells were intact) were studied in 5 to 10 cells per animal. From each animal, 150 cells not introduced to fura 2-AM and without morphological alteration were measured for length and midpoint width. Cell volume was estimated as cell length  $\times$  width  $\times 0.00759$ , as established by 2D light and 3D confocal microscopy.<sup>19</sup>

### Vascular Function

L-shaped holders were inserted into the lumen of 2- to 4-mm segments of the common carotid arteries; one holder was connected to a force-displacement transducer and the other to a micrometer in organ baths containing Krebs buffer and indomethacin.<sup>20</sup> After gradually increasing tension to 1000 mg and exposure to 60 mmol/L  $K^+$ ,  $3 \times 10^{-7}$  mol/L phenylephrine, and  $10^{-4}$  mol/L acetylcholine to ensure reactivity, segments were equilibrated 30 minutes before experiments started. Four segments from each animal were precontracted with phenylephrine ( $3 \times 10^{-7}$  mol/L) and relaxed with cumulative doses of acetylcholine (2 segments) and  $Na^+$  nitroprusside (1 segment), whereas 1 segment was also pretreated with  $10^{-4}$  mol/L  $N^G$ -nitro-L-arginine methyl ester (L-NAME) before exposure to acetylcholine.

### Allometric Scaling

In addition to exercise, differences in cardiac muscle weight and  $\dot{V}O_2$  may result from altered body mass.<sup>21</sup> According to dimensional analysis and empirical studies,  $\dot{V}O_2$  should be expressed in relation to body mass raised to the power of 0.75,<sup>22</sup> whereas ventricular mass should be expressed with the scaling exponent 0.78, which empirically is the best approximation when lean body mass is unavailable.<sup>23</sup>

### Statistics


Data are expressed as mean  $\pm$  SD.  $EC_{50}$  values were obtained as previously described.<sup>24</sup> The Friedman test, Wilcoxon paired samples  $t$  test, and appropriate procedures for multiple comparisons determined changes in  $\dot{V}O_{2\max}$ , whereas the Mann-Whitney test determined different cellular data, cardiac weights, and arterial function; a univariate repeated-measures ANOVA including Scheffé post hoc tests verified the differences. Relationships were assessed by Pearson's correlation coefficient and complementary univariate, forward and backward linear regression analyses.  $\dot{V}O_{2\max}$  was modeled using cardiomyocyte volume, length, width, fractional shortening, time to half contraction and relaxation, and time to half  $Ca^{2+}$  peak and decay, and vasorelaxation  $EC_{50}$  for detraining as explanatory variables, with  $P > 0.05$  as exclusion criterion.

## Results

Both exercise training and detraining led to substantial increase and regression of aerobic fitness and  $\dot{V}O_{2\max}$ , which closely corresponded to changes in cardiomyocyte contractile capacity,  $Ca^{2+}$  handling, and arterial endothelium function.

### $\dot{V}O_{2\max}$ During Training and Detraining

As previously reported,<sup>6,18</sup> regular high-intensity interval running increased  $\dot{V}O_{2\max}$  substantially. After 8 to 10 weeks




**Figure 2.** Relationship between  $\dot{V}O_{2max}$  and cardiomyocyte volume (A), fractional shortening (B), half-time diastolic relaxation (C), and half-time  $\text{Ca}^{2+}$  decay (D) and between cardiomyocyte fractional shortening and cell volume (E) in rats trained for 2, 4, 8, and 13 weeks and sedentary controls, respectively. Individual data.<sup>11,21</sup>

of training,  $\dot{V}O_{2max}$  stabilized 37% above baseline and 26% above sedentary. During detraining, exercise-gained  $\dot{V}O_{2max}$  decreased 50% within 2 weeks and stabilized 5% above sedentary after 4 weeks (Figure 1). In trained animals, univariate analysis of the data revealed that cardiomyocyte length, relaxation, shortening,  $\text{Ca}^{2+}$  decay, and volume correlated strongly with  $\dot{V}O_{2max}$  (Figure 2). Backward multiple regression identified cell length, diastolic relaxation, and  $\text{Ca}^{2+}$  decay as the main factors for  $\dot{V}O_{2max}$ ; unstandardized coefficients  $b$   $0.95 \pm \text{SE } 0.39$ ,  $P < 0.01$ ;  $-0.71 \pm 0.54$ ,  $P < 0.01$ ; and  $-1.02 \pm 0.42$ ,  $P < 0.02$ , respectively; residual SD=4.84, adjusted  $R^2=0.87$ . In detraining, cell hypertrophy regression correlated closely with  $\dot{V}O_{2max}$ . Strong correlation also occurred between detraining-induced regressed  $\dot{V}O_{2max}$  and cardiomyocyte shortening and  $\text{Ca}^{2+}$  decay, whereas a trend

occurred for myocyte relaxation (Figure 3). Acetylcholine-induced relaxation correlated less markedly with  $\dot{V}O_{2max}$  ( $r = -0.37$ ,  $P < 0.05$ ) than cardiomyocyte variables. Backward multiple regression identified cardiomyocyte length and endothelium-dependent arterial relaxation as the main determinants for changes in  $\dot{V}O_{2max}$  during detraining with unstandardized coefficients  $b$   $0.80 \pm 0.09$ ,  $P < 0.01$ , and  $-4.67 \pm 1.85$ ,  $P < 0.02$ , respectively; residual SD=4.63, adjusted  $R^2=0.76$ .

### Cardiomyocyte Morphology and Function

Training increased ventricular weights and cardiomyocyte dimensions and improved contractility and  $\text{Ca}^{2+}$  handling in the heart<sup>6,18</sup> (Figure 1). Animals randomized for detraining also increased cardiomyocyte width and length by 20% to 22% and estimated volume by 46% (Figure 4). During



**Figure 3.** Relationship between  $\dot{V}O_{2\max}$  and cardiomyocyte volume (A), fractional shortening (B), half-time diastolic relaxation (C), and half-time  $Ca^{2+}$  decay (D) in detrained and sedentary rats. Training lasted 10 weeks (TR10) and detraining 2 (DETR2) and 4 (DETR4) weeks, respectively. Individual data.

detraining, the responses varied slightly; cell width regressed completely within 2 weeks, whereas length (7% to 5%) and volume (15% to 9%) remained enlarged after 2 and 4 weeks, respectively, ie, similar to  $\dot{V}O_{2\max}$ . Parallel changes occurred in cardiac weights, which regressed toward sedentary within 4 weeks of detraining (Table).

Cardiomyocyte  $Ca^{2+}$  handling and intrinsic contractility were assessed at physiological cell stimulation frequencies (Figure 5). Within 2 weeks of detraining, the  $\approx 30\%$  increase in cardiomyocyte fractional shortening regressed almost completely. Diastolic and systolic  $fura\ 2\ Ca^{2+}$  ratios and amplitude of  $Ca^{2+}$  transient were largely unaffected by training/detraining. The increase in cardiomyocyte contractility seemed to be associated with higher myofilament  $Ca^{2+}$  sensitivity. The  $Ca^{2+}$  sensitivity index (cell shortening/ $Ca^{2+}$  ratio amplitude) was elevated at 7 to 10 Hz electrical stimulation after 10 weeks of training and reversed to sedentary values during detraining. As shown in Figure 6, rates of both contraction and relaxation were increased, with parallel changes in  $Ca^{2+}$  handling. These changes prevailed to some extent after 2 weeks of detraining but not by week 4.

### Endothelium-Dependent Arterial Relaxation

Endothelium-dependent arterial relaxation increased significantly after regular exercise training. After 10 weeks, the magnitude of acetylcholine-induced relaxation increased by 13% and  $EC_{50}$  for agonist decreased 4-fold, whereas maximal absolute relaxation ( $R_{\max}$ ) leveled off 24% above sedentary (Figure 7). With detraining, all


effects reversed within 2 weeks. The 7% and 2-fold larger  $EC_{50}$  and 11% increased  $R_{\max}$  after cumulative  $Na^+$  nitroprusside addition in trained animals indicate a transient enhanced sensitivity to nitric oxide (NO), because it vanished within 2 weeks of detraining.

### Discussion

The present training-detraining experiments identified 2 distinctive cellular factors associated with changes in aerobic fitness, one closely correlated with cardiac myocyte size and function and another related to endothelium-dependent arterial relaxation. Although both may be important for the salutary health effects of exercise, the myocardial mechanisms seem to be more closely correlated to  $\dot{V}O_{2\max}$ .

### Cardiomyocyte Contractile Capacity

During long-term adaptation to regular exercise, the heart meets increased needs of peripheral tissues by matching pump capacity to afford sufficient cardiac output to transport oxygen corresponding to  $\dot{V}O_{2\max}$ .<sup>25</sup> Both physiological hypertrophy and changes in myocardial function may account for the required increase in stroke volume. In the present study, changes in  $\dot{V}O_{2\max}$  closely paralleled cardiomyocyte length and width, providing a cellular basis for increase and regression of right and left ventricle stroke volumes. In addition, long-term changes in myocardial function may contribute significantly by altering diastolic filling and systolic emptying,<sup>12,14,26</sup> as indicated by higher



**Figure 4.** Cardiomyocyte length (A), width (B), and estimated volume (C) in detrained and sedentary rats (SED10, SED2, SED4) in 4800 cells, 150 from each rat. Training lasted 10 weeks (TR10) and detraining 2 (DETR2) and 4 (DETR4) weeks, respectively. Data are mean±SD. TR/DETR vs sedentary: \* $P < 0.01$ , † $P < 0.05$ . TR10 vs DETR2: ‡ $P < 0.01$ , § $P < 0.05$ , and DETR4: || $P < 0.01$ .

ejection fraction and fraction of shortening after training.<sup>6,18,27,28</sup> Several aspects of cardiomyocyte contractile function corresponded to training- and detraining-induced changes in  $\dot{V}O_2\text{max}$ , including cardiomyocyte shortening and relaxation representing systolic and diastolic contractile properties, respectively. Although stroke volume is acutely regulated by extracardiac factors such as venous

**Body Mass and Cardiac Weights**

	Sedentary	TR10	DETR2	DETR4
Body mass, g				
Before	239.4±11.0	232.8±12.5‡	254.5±12.9	243.2±15.2
After	303.9±15.3	288.0±27.1‡	325.5±19.3	309.5±27.1
Heart weight				
In mg	1422.8±140.3	1569.2±150.6	1697.2±55.3§	1482.7±186.3
In mg/g	4.7±0.4	5.5±0.3*	5.2±0.3	4.8±0.8
In mg/g <sup>0.78</sup>	16.5±1.4	18.9±1.1†	18.6±0.9†	17.1±2.8
LV weight				
In mg	1051.3±109.7	1162.5±126.6	1263.1±80.6†	1111.9±172.5
In mg/g	3.5±0.3	4.0±0.2*	3.9±0.4	3.6±0.7
In mg/g <sup>0.78</sup>	12.2±1.1	14.0±0.8†	13.9±1.4	12.8±2.4
RV weight				
In mg	371.5±52.6	406.8±61.8	434.1±81.0	370.8±40.0
In mg/g	1.2±0.2	1.4±0.2	1.3±0.2	1.2±0.2
In mg/g <sup>0.78</sup>	4.3±0.6	4.9±0.8	4.7±0.7	4.3±0.6


Body mass before and after experimental period and postmortem heart weights in detrained and sedentary; heart weights after Langendorff perfusion (see Methods). Training lasted 10 weeks (TR10); detraining, 2 (DETR2) and 4 (DETR4) weeks, respectively. LV indicates left ventricle; RV, right ventricle. Data are mean±SD.

TR/DETR vs sedentary: \* $P < 0.01$ , † $P < 0.05$ .  
 TR10 vs DETR2: ‡ $P < 0.05$ .  
 DETR2 vs DETR4: § $P < 0.05$ .

return, neurohormonal regulation, and afterload, previous studies identified significant correlation between contractile function of isolated myocytes and integrated in vivo function, indicating a contribution of intrinsic myocardial properties as well at both the molecular<sup>29</sup> and cellular levels.<sup>6,18,30</sup> Both statistical correlation and parallel time courses indicate that changes in cardiomyocyte size and function are likely to account for the changes in  $\dot{V}O_2\text{max}$ . Because of the high internal correlations between different measures of cardiomyocyte size and function, it is to be expected that only 1 or 2 prove to be significant in multivariate regression analysis. However, this statistical interdependence does not preclude the possibility that myocyte size, contractility, and relaxation contribute cumulatively to the total contractile capacity in vivo.

**Endothelium-Dependent Arterial Relaxation**


The training-detraining experiment demonstrated substantial changes in acetylcholine-induced arterial relaxation, which were highly sensitive to inhibition of the endothelial nitric oxide synthase (eNOS) inhibitor L-NAME. These observations indicate induction and regression of endothelial function, which were significantly correlated with  $\dot{V}O_2\text{max}$  in both univariate and multivariate analysis. However, the marked difference in time course of  $EC_{50}$  for acetylcholine indicates that endothelium-dependent vasodilation may not be as directly related to  $\dot{V}O_2\text{max}$  as cardiomyocyte size and function. Whereas training-induced myocardial effects gradually regress over 3 to 4 weeks, exercise-gained endothelium-dependent relaxation is completely abolished within 2 weeks. The time course of the onset of endothelium-dependent changes could not be




**Figure 5.** Cardiomyocyte contractility and Ca<sup>2+</sup> handling in detrained and sedentary rats determined in 4 to 10 cells per rat. Training lasted 10 weeks (TR10) and detraining 2 (DETR2) and 4 (DETR4) weeks, respectively. A, Cell shortening; B, Ca<sup>2+</sup> ratio amplitude; C, Ca<sup>2+</sup> ratio sensitivity index (cardiomyocyte relative shortening/Ca<sup>2+</sup> ratio amplitude); and D, diastolic and systolic Ca<sup>2+</sup> ratios. Data are mean±SD. TR10 vs sedentary: \**P*<0.05; TR10 vs DETR2: †*P*<0.05; and DETR4: ‡*P*<0.01, §*P*<0.05.

determined because the previous studies did not include arterial function. On the basis of pilot experiments and exercise-induced resistance to decompression,<sup>31</sup> we hypothesize that endothelial function changes much more quickly than myocardial. A more rapid time course does

not preclude salutary health effects. Hambrecht et al<sup>7</sup> recently demonstrated training-induced improvement in myocardial oxygen supply associated with increased endothelium-dependent relaxation and upregulation of the eNOS signaling pathway.



**Figure 6.** Time course of contraction/relaxation and Ca<sup>2+</sup> transient in detrained and sedentary cardiomyocytes stimulated at increasing frequencies. Training lasted 10 weeks (TR10) and detraining 2 (DETR2) and 4 (DETR4) weeks, respectively. A and B, Time to peak contraction and peak Ca<sup>2+</sup> ratio, respectively; C and D, half-time to peak contraction and peak Ca<sup>2+</sup> ratio, respectively; E and F, half-time to relaxation and Ca<sup>2+</sup> ratio decay, respectively. Data are mean±SD. TR10 vs sedentary: \**P*<0.01, †*P*<0.05. DETR2 vs sedentary: ‡*P*<0.05. TR10 vs DETR2: §*P*<0.01, ||*P*<0.05, and DETR4: #*P*<0.01.



**Figure 7.** Phenylephrine-precontracted carotid artery response to accumulating acetylcholine, acetylcholine+L-NAME, or Na<sup>+</sup> nitroprusside, all in presence of indomethacin, in trained 10 weeks (TR10, A), detrained 2 weeks (DETR2, B), and detrained 4 weeks (DETR4, C), with respective controls. Dose-response curves are constructed as described in Reference 27. TR10 vs sedentary: \*P<0.01, †P<0.05. Note that training-induced responses were lost within 2 weeks of detraining.

**Conclusions**

Regular exercise induces substantial improvements in cardiomyocyte and endothelial function that are likely to contribute significantly to improved health and increased resistance to cardiovascular disease. The present study demonstrates that both  $\dot{V}O_2\text{max}$  and myocardial effects plateau within 6 to 8 weeks and regress almost completely within 4 weeks of

detraining and suggests an even more rapid time course for endothelium-dependent arterial relaxation. Although both myocardial and endothelium-dependent factors correlate significantly with  $\dot{V}O_2\text{max}$ , the parallel temporal relationship of cardiomyocyte hypertrophy and contractile function indicate that myocardial cellular mechanisms may be more important for increased aerobic capacity. Studies in progress will more accurately define the time course of exercise-induced changes in endothelial function and determine whether training intensity affects magnitude of myocardial and endothelial responses differently and how these findings apply in heart failure.

**Acknowledgments**

This study was supported by grants from the National Council on Cardiovascular Diseases, the St Olav’s Hospital, and the Torstein Erbo, Arild and Emilie Bachke, Ingeborg and Anders Nordheim, and Randi and Hans Arnet Foundations. Dr Kemi is the recipient of a research fellowship from the Norwegian University of Science and Technology. We thank Ragnhild Støen for introducing us to arterial relaxation experiments.

**References**

1. Belardinelli R, Georgiou D, Cianci G, et al. Randomized, controlled trial of long-term moderate exercise training in chronic heart failure: effects on functional capacity, quality of life, and clinical outcome. *Circulation*. 1999;99:1173–1182.
2. Myers J, Prakash M, Froelicher V, et al. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med*. 2002;346:793–801.
3. Frenzel H, Schwartzkopff B, Holtermann W, et al. Regression of cardiac hypertrophy: morphometric and biochemical studies in rat heart after swimming training. *J Mol Cell Cardiol*. 1988;20:737–751.
4. Kemi OJ, Loennechen JP, Wisløff U, et al. Intensity-controlled treadmill running in mice: cardiac and skeletal muscle hypertrophy. *J Appl Physiol*. 2002;93:1301–1309.
5. Wisløff U, Helgerud J, Kemi OJ, et al. Intensity-controlled treadmill running in rats:  $\dot{V}O_2\text{max}$  and cardiac hypertrophy. *Am J Physiol*. 2001;280:H1301–H1310.
6. Wisløff U, Loennechen JP, Falck G, et al. Increased contractility and calcium sensitivity in cardiac myocytes isolated from endurance trained rats. *Cardiovasc Res*. 2001;50:495–508.
7. Hambrecht R, Adams V, Erbs S, et al. Regular physical activity improves endothelial function in patients with coronary artery disease by increasing phosphorylation of endothelial nitric oxide synthase. *Circulation*. 2003;107:3152–3158.
8. Hambrecht R, Fiehn E, Weigl C, et al. Regular physical exercise corrects endothelial dysfunction and improves exercise capacity in patients with chronic heart failure. *Circulation*. 1998;98:2709–2715.
9. Arvola P, Wu X, Kähönen M, et al. Exercise enhances vasorelaxation in experimental obesity associated hypertension. *Cardiovasc Res*. 1999;43:992–1002.
10. Minami A, Ishimura N, Harada N, et al. Exercise training improves acetylcholine-induced endothelium-dependent hyperpolarization in type 2 diabetic rats, Otsuka Long-Evans Tokushima fatty rats. *Atherosclerosis*. 2002;162:85–92.
11. Mujika I, Padilla S. Cardiorespiratory and metabolic characteristics of detraining in humans. *Med Sci Sports Exerc*. 2001;33:413–421.
12. Pelliccia A, Maron BJ, De Luca R, et al. Remodeling of left ventricular hypertrophy in elite athletes after long-term deconditioning. *Circulation*. 2002;105:944–949.
13. Cullinane EM, Sady SP, Vadebonceur L, et al. Cardiac size and  $\dot{V}O_2\text{max}$  do not decrease after short-term exercise cessation. *Med Sci Sports Exerc*. 1986;18:420–424.
14. Martin WH III, Coyle EF, Bloomfield SA, et al. Effects of physical deconditioning after intense endurance training on left ventricular dimensions and stroke volume. *J Am Coll Cardiol*. 1986;7:982–989.
15. Craig BW, Martin G, Betts J, et al. The influence of training-detraining upon the heart, muscle and adipose tissue of female rats. *Mech Ageing Dev*. 1991;57:49–61.

16. Hickson RC, Galassi TM, Dougherty KA. Repeated development and regression of exercise-induced cardiac hypertrophy in rats. *J Appl Physiol*. 1983;54:794–797.
17. Hickson RC, Hammons GT, Holoszy JO. Development and regression of exercise-induced cardiac hypertrophy in rats. *Am J Physiol*. 1979;236:H268–H272.
18. Wisløff U, Loennechen JP, Currie S, et al. Aerobic exercise reduces cardiomyocyte hypertrophy and increases contractility, Ca<sup>2+</sup> sensitivity and SERCA-2 in rat after myocardial infarction. *Cardiovasc Res*. 2002;54:162–174.
19. Satoh H, Delbridge LMD, Blatter LA, et al. Surface:volume relationship in cardiac myocytes studied with confocal microscopy and membrane capacitance measurements: species-dependence and developmental effects. *Biophys J*. 1996;70:1494–1504.
20. Støen R, Brubakk AM, Vik T, et al. Postnatal changes in mechanisms mediating acetylcholine-induced relaxation in piglet femoral arteries. *Pediatr Res*. 1997;41:702–707.
21. Darveau CA, Suarez RK, Andrews RD, et al. Allometric cascade as a unifying principle of body mass effects on metabolism. *Nature*. 2002;417:166–170.
22. Taylor CR, Maloiy GM, Weibel EM, et al. Design of the mammalian respiratory system, III: scaling maximum aerobic capacity to body mass: wild and domestic mammals. *Respir Physiol*. 1981;44:25–37.
23. Batterham AM, George KP, Mullineaux DR. Allometric scaling of left ventricular mass by body dimensions in males and females. *Med Sci Sports Exerc*. 1997;29:181–186.
24. Ariëns EJ, Simonis AM, van Rossum JM. Drug-receptor interactions: interaction of one or more drugs with one receptor system. In: Ariëns EJ, ed. *Molecular Pharmacology*. New York, NY: Academic; 1964: 119–286.
25. Wagner PD. A theoretical analysis of factors determining Vo<sub>2</sub>MAX at sea level and altitude. *Respir Physiol*. 1996;106:329–343.
26. Gledhill N, Cox D, Jamnik R. Endurance athletes' stroke volume does not plateau: major advantage is diastolic function. *Med Sci Sports Exerc*. 1994;26:1116–1121.
27. Moore RL, Musch TI, Yelamarty RV, et al. Chronic exercise alters contractility and morphology of isolated rat cardiac myocytes. *Am J Physiol*. 1993;264:C1180–C1189.
28. Wisløff U, Helgerud J, Støylen A, et al. Atrioventricular plane displacement in female endurance athletes. *Med Sci Sports Exerc*. 2001;33: 1503–1510.
29. Loennechen JP, Støylen A, Beisvag V, et al. Regional expression of endothelin-1, ANP, IGF-1, and LV wall stress in the infarcted rat heart. *Am J Physiol*. 2001;280:H2902–H2910.
30. Loennechen JP, Wisløff U, Falck G, et al. Cardiomyocyte contractility and calcium handling partially recover after early deterioration during post-infarction failure in rat. *Acta Physiol Scand*. 2002;176: 17–26.
31. Wisløff U, Richardson RS, Brubakk AO. NOS inhibition increases bubble formation and reduces survival in sedentary but not exercised rats. *J Physiol*. 2003;15:577–582.

## Aerobic Fitness Is Associated With Cardiomyocyte Contractile Capacity and Endothelial Function in Exercise Training and Detraining

Ole Johan Kemi, Per Magnus Haram, Ulrik Wisløff and Øyvind Ellingsen

*Circulation*. 2004;109:2897-2904; originally published online June 1, 2004;

doi: 10.1161/01.CIR.0000129308.04757.72

*Circulation* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2004 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circ.ahajournals.org/content/109/23/2897>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation* is online at:  
<http://circ.ahajournals.org/subscriptions/>