

Part 7: CPR Techniques and Devices

2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care

Diana M. Cave, Chair; Raul J. Gazmuri; Charles W. Otto; Vinay M. Nadkarni; Adam Cheng; Steven C. Brooks; Mohamud Daya; Robert M. Sutton; Richard Branson; Mary Fran Hazinski

Over the past 25 years a variety of alternatives to conventional manual CPR have been developed in an effort to enhance perfusion during attempted resuscitation from cardiac arrest and to improve survival. Compared with conventional CPR, these techniques and devices typically require more personnel, training, and equipment, or they apply to a specific setting. Application of these devices has the potential to delay or interrupt CPR, so rescuers should be trained to minimize any interruption of chest compressions or defibrillation and should be retrained as needed. Efficacy for some techniques and devices has been reported in selected settings and patient conditions; however, no alternative technique or device in routine use has consistently been shown to be superior to conventional CPR for out-of-hospital basic life support. In this section, no class of recommendation is made when there is insufficient evidence of benefit or harm, particularly if human data are extremely limited. For those devices assigned a 2005 Class of Recommendation other than Indeterminate, Classes of Recommendation were assigned when possible using the same criteria applied throughout this document (see Part 1: “Executive Summary” and Part 2: “Evidence Evaluation”).

Whenever these devices are used, providers should monitor for evidence of benefit versus harm. The experts are aware of several clinical trials of the devices listed below that are under way and/or recently concluded, so readers are encouraged to monitor for the publication of additional trial results in peer-reviewed journals and AHA scientific advisory statements.

CPR Techniques

High-Frequency Chest Compressions

High-frequency chest compression (typically at a frequency >120 per minute) has been studied as a technique for improving resuscitation from cardiac arrest.¹ The sparse human data have demonstrated mixed results. One clinical trial including 9 patients² and another including 23 patients³ showed that a compression frequency of 120 per minute improved hemodynamics compared to conven-

tional chest compressions; no change in clinical outcome was reported. These 2010 AHA Guidelines for CPR and ECC recommend compressions at a rate of at least 100/min. There is insufficient evidence to recommend the routine use of high-frequency chest compressions for cardiac arrest. However, high-frequency chest compressions may be considered by adequately trained rescue personnel as an alternative (Class IIB, LOE C).

Open-Chest CPR

In open-chest CPR the heart is accessed through a thoracotomy (typically created through the 5th left intercostal space) and compression is performed using the thumb and fingers, or with the palm and extended fingers against the sternum. Use of this technique generates forward blood flow and coronary perfusion pressure that typically exceed those generated by closed chest compressions.

There are few human studies comparing open-chest CPR to conventional CPR in cardiac arrest and no prospective randomized trials. Several studies of open-chest CPR have demonstrated improved coronary perfusion pressure and/or return of spontaneous circulation (ROSC) for both the in-hospital (eg, following cardiac surgery)^{4–6} and out-of-hospital environments.^{7–10}

Several small case series of cardiac arrest patients treated with thoracotomy and open-chest CPR after blunt^{11,12} or penetrating trauma^{12–14} reported survivors with mild or no neurological deficit.

There is insufficient evidence of benefit or harm to recommend the routine use of open-chest CPR. However, open-chest CPR can be useful if cardiac arrest develops during surgery when the chest or abdomen is already open, or in the early postoperative period after cardiothoracic surgery (Class IIA, LOE C). A resuscitative thoracotomy to facilitate open-chest CPR may be considered in very select circumstances of adults and children with out-of-hospital cardiac arrest from penetrating trauma with short transport times to a trauma facility (Class IIB, LOE C).^{15,16}

The American Heart Association requests that this document be cited as follows: Cave DM, Gazmuri RJ, Otto CW, Nadkarni VM, Cheng A, Brooks SC, Daya M, Sutton RM, Branson R, Hazinski MF. Part 7: CPR techniques and devices: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2010;122(suppl 3):S720–S728.

(*Circulation*. 2010;122[suppl 3]:S720–S728.)

© 2010 American Heart Association, Inc.

Circulation is available at <http://circ.ahajournals.org>

DOI: 10.1161/CIRCULATIONAHA.110.970970

Interposed Abdominal Compression-CPR

The interposed abdominal compression (IAC)-CPR is a 3-rescuer technique (an abdominal compressor plus the chest compressor and the rescuer providing ventilations) that includes conventional chest compressions combined with alternating abdominal compressions. The dedicated rescuer who provides manual abdominal compressions will compress the abdomen midway between the xiphoid and the umbilicus during the relaxation phase of chest compression. Hand position, depth, rhythm, and rate of abdominal compressions are similar to those for chest compressions and the force required is similar to that used to palpate the abdominal aorta. In most reports, an endotracheal tube is placed before or shortly after initiation of IAC-CPR. IAC-CPR increases diastolic aortic pressure and venous return, resulting in improved coronary perfusion pressure and blood flow to other vital organs.

In 2 randomized in-hospital trials, IAC-CPR performed by trained rescuers improved short-term survival¹⁷ and survival to hospital discharge¹⁸ compared with conventional CPR for adult cardiac arrest. The data from these studies were combined in 2 positive meta-analyses.^{19,20} However, 1 randomized controlled trial of adult out-of-hospital cardiac arrest²¹ did not show any survival advantage to IAC-CPR. Although there were no complications reported in adults,¹⁹ 1 pediatric case report²² documented traumatic pancreatitis following IAC-CPR.

IAC-CPR may be considered during in-hospital resuscitation when sufficient personnel trained in its use are available (Class IIb, LOE B). There is insufficient evidence to recommend for or against the use of IAC-CPR in the out-of-hospital setting or in children.

“Cough” CPR

“Cough” CPR describes the use of forceful voluntary coughs every 1 to 3 seconds in conscious patients shortly after the onset of a witnessed nonperfusing cardiac rhythm in a controlled environment such as the cardiac catheterization laboratory. Coughing episodically increases the intrathoracic pressure and can generate systemic blood pressures higher than those usually generated by conventional chest compressions,^{23,24} allowing patients to maintain consciousness^{23–26} for a brief arrhythmic interval (up to 92 seconds documented in humans).²⁵

“Cough” CPR has been reported exclusively in awake, monitored patients (predominantly in the cardiac catheterization laboratory) when arrhythmic cardiac arrest can be anticipated, the patient remains conscious and can be instructed before and coached during the event, and cardiac activity can be promptly restored.^{23–33} However, not all victims are able to produce hemodynamically effective coughs.²⁷

“Cough” CPR is not useful for unresponsive victims and should not be taught to lay rescuers. “Cough” CPR may be considered in settings such as the cardiac catheterization laboratory for conscious, supine, and monitored patients if the patient can be instructed and coached to cough forcefully every 1 to 3 seconds during the initial seconds of an arrhythmic cardiac arrest. It should not delay definitive treatment (Class IIb, LOE C).

Prone CPR

When the patient cannot be placed in the supine position, it may be reasonable for rescuers to provide CPR with the patient in the prone position, particularly in hospitalized patients with an advanced airway in place (Class IIb, LOE C).^{34–37}

Precordial Thump

This section is new to the 2010 Guidelines and is based on the conclusions reached by the 2010 ILCOR evidence evaluation process.³⁸

A precordial thump has been reported to convert ventricular tachyarrhythmias in 1 study with concurrent controls,³⁹ single-patient case reports, and small case series.^{40–44} However, 2 larger case series found that the precordial thump was ineffective in 79 (98.8%) of 80 cases⁴⁵ and in 153 (98.7%) of 155 cases of malignant ventricular arrhythmias.⁴⁶ Case reports and case series^{47–49} have documented complications associated with precordial thump including sternal fracture, osteomyelitis, stroke, and triggering of malignant arrhythmias in adults and children.

The precordial thump should not be used for unwitnessed out-of-hospital cardiac arrest (Class III, LOE C). The precordial thump may be considered for patients with witnessed, monitored, unstable ventricular tachycardia including pulseless VT if a defibrillator is not immediately ready for use (Class IIb, LOE C), but it should not delay CPR and shock delivery. There is insufficient evidence to recommend for or against the use of the precordial thump for witnessed onset of asystole.

Percussion Pacing

Percussion (eg, fist) pacing refers to the use of regular, rhythmic and forceful percussion of the chest with the rescuer’s fist in an attempt to pace the myocardium. There is little evidence supporting fist or percussion pacing in cardiac arrest based on 6 single-patient case reports^{50–55} and a moderate-sized case series.⁵⁶ There is insufficient evidence to recommend percussion pacing during typical attempted resuscitation from cardiac arrest.

CPR Devices

Devices to Assist Ventilation

Automatic and Mechanical Transport Ventilators

Automatic Transport Ventilators

There are very few studies evaluating the use of automatic transport ventilators (ATVs) during attempted resuscitation in patients with endotracheal intubation. During prolonged resuscitation efforts, the use of an ATV (pneumatically powered and time- or pressure-cycled) may provide ventilation and oxygenation similar to that possible with the use of a manual resuscitation bag, while allowing the Emergency Medical Services (EMS) team to perform other tasks (Class IIb, LOE C^{57,58}). Disadvantages of ATVs include the need for an oxygen source and a power source. Thus, providers should always have a bag-mask device available for manual backup.

For additional information regarding support of airway and ventilation in the adult, see ACLS Part 8.1 in these Guidelines.

Manually Triggered, Oxygen-Powered, Flow-Limited Resuscitators

In a study of 104 anesthetized nonarrest patients without an advanced airway in place (ie, no endotracheal tube; patients were ventilated through a mask), patients ventilated by firefighters with manually triggered, oxygen-powered, flow-limited resuscitators had less gastric inflation than those ventilated with a bag-mask device.⁵⁹ Manually triggered, oxygen-powered, flow-limited resuscitators may be considered for the management of patients who do not have an advanced airway in place and for whom a mask is being used for ventilation during CPR (Class IIb, LOE C). Rescuers should avoid using the automatic mode of the oxygen-powered, flow-limited resuscitator during CPR because it may generate high positive end-expiratory pressure (PEEP) that may impede venous return during chest compressions and compromise forward blood flow (Class III, LOE C⁶⁰).

Devices to Support Circulation

Active Compression-Decompression CPR

Active compression-decompression CPR (ACD-CPR) is performed with a device that includes a suction cup to actively lift the anterior chest during decompression. The application of external negative suction during the decompression phase of CPR creates negative intrathoracic pressure and thus potentially enhances venous return to the heart. When used, the device is positioned at midsternum on the chest.

Results from the use of ACD-CPR have been mixed. In several studies^{61–66} ACD-CPR improved ROSC and short-term survival compared with conventional CPR. Of these studies, 3 showed improvement in neurologically intact survival.^{61,64,65} In contrast, 1 Cochrane meta-analysis of 10 studies involving both in-hospital arrest (826 patients) and out-of-hospital arrest (4162 patients)⁶⁷ and several other controlled trials^{68–74} comparing ACD-CPR to conventional CPR showed no difference in ROSC or survival. The meta-analysis⁶⁷ did not find any increase in ACD-CPR-related complications.

There is insufficient evidence to recommend for or against the routine use of ACD-CPR. ACD-CPR may be considered for use when providers are adequately trained and monitored (Class IIb, LOE B).

Phased Thoracic-Abdominal Compression-Decompression CPR With a Handheld Device

Phased thoracic-abdominal compression-decompression CPR (PTACD-CPR) combines the concepts of IAC-CPR and ACD-CPR. A handheld device alternates chest compression and abdominal decompression with chest decompression and abdominal compression. Evidence from 1 prospective randomized clinical study of adults in cardiac arrest⁷⁵ demonstrated no improvement in survival to hospital discharge with use of PTACD-CPR during out-of-hospital cardiac arrest.

There is insufficient evidence to support or refute the use of PTACD-CPR for the treatment of cardiac arrest.

Impedance Threshold Device

The impedance threshold device (ITD) is a pressure-sensitive valve that is attached to an endotracheal tube, supraglottic airway, or face mask. The ITD limits air entry into the lungs during the decompression phase of CPR, creating negative intrathoracic pressure and improving venous return to the heart and cardiac output during CPR. It does so without impeding positive pressure ventilation or passive exhalation.

Originally, the ITD was used with a cuffed endotracheal tube during bag-tube ventilation and ACD-CPR.^{76–78} The ITD and ACD-CPR devices are thought to act synergistically to enhance venous return. During ACD-CPR with or without the ITD, 1 randomized study⁷⁶ found no difference in survival, whereas another randomized study⁷⁹ found that the addition of an ITD improved short-term survival (24-hour survival and survival to ICU admission).

The ITD also has been used during conventional CPR with an endotracheal tube or with a face mask, if a tight seal is maintained.^{77,80,81} During conventional CPR with and without the ITD, 1 randomized trial⁸⁰ reported no difference in overall survival; however, 1 prospective cohort study⁸² reported improved survival to emergency department (ED) admission with the use of the ITD. One meta-analysis of pooled data from both conventional CPR and ACD-CPR randomized trials⁸³ demonstrated improved ROSC and short-term survival associated with the use of an ITD in the management of adult out-of-hospital cardiac arrest patients but no significant improvement in either survival to hospital discharge or neurologically intact survival to discharge.

Three cohort studies with historic controls that implemented 2005 Guidelines plus ITD demonstrated improved survival to hospital discharge for out-of-hospital cardiac arrest.^{84–86} It was not possible to determine the relative contribution of the ITD to the improved outcome. The use of the ITD may be considered by trained personnel as a CPR adjunct in adult cardiac arrest (Class IIb, LOE B).

Mechanical Piston Devices

A mechanical piston device consists of a compressed gas- or electric-powered plunger mounted on a backboard; it is used to depress the sternum. Some incorporate a suction cup in the piston device while others do not. In 3 studies^{87–89} the use of a mechanical piston device for CPR improved end-tidal CO₂ and mean arterial pressure during adult cardiac arrest resuscitation. However, compared with manual CPR, no improvement in short- and long-term survival in adult patients was demonstrated.^{87,90} Initiation and removal of the mechanical piston device were noted to increase interruptions in CPR.⁹¹

The Lund University Cardiac Arrest System (LUCAS) is a gas- (oxygen or air) or electric-powered piston device that produces a consistent chest compression rate and depth. It incorporates a suction cup attached to the sternum that returns the sternum to the starting position. There are no randomized control trials comparing the device with conventional CPR in human cardiac arrests. One case

series with concurrent controls⁹² showed no benefit over conventional CPR for out-of-hospital witnessed cardiac arrest. Additional case series have reported variable success with the device.^{93–98} One feasibility study reported successful deployment during diagnostic and interventional procedures.⁹⁹

There is insufficient evidence to support or refute the routine use of mechanical piston devices in the treatment of cardiac arrest. Mechanical piston devices may be considered for use by properly trained personnel in specific settings for the treatment of adult cardiac arrest in circumstances (eg, during diagnostic and interventional procedures) that make manual resuscitation difficult (Class IIb, LOE C). Rescuers should attempt to limit substantial interruptions in CPR during deployment. The device should be programmed to deliver high-quality CPR, ensuring an adequate compression depth of at least 2 inches (5 cm)—this may require conversion from a percent of chest depth, a rate of at least 100 compressions per minute, and a compression duration of approximately 50% of the cycle length.

Load-Distributing Band CPR or Vest CPR

The load-distributing band (LDB) is a circumferential chest compression device composed of a pneumatically or electrically actuated constricting band and backboard. Case series have demonstrated improved hemodynamics,¹⁰⁰ ROSC,^{101,102} and survival to hospital discharge with use of the LDB for cardiac arrest.¹⁰² In a study using concurrent controls,¹⁰³ the use of LDB-CPR was associated with lower odds of 30-day survival (odds ratio 0.4). One multicenter prospective randomized controlled trial^{104,104A} comparing LDB-CPR (Auto-pulse device) to manual CPR for out-of-hospital cardiac arrest demonstrated no improvement in 4-hour survival and worse neurologic outcome when the device was used. These results raised concerns about possible harm with use of this device. Further studies are required to determine whether site-specific factors¹⁰⁵ and experience with deployment of the device¹⁰⁶ could influence its efficacy.

The LDB may be considered for use by properly trained personnel in specific settings for the treatment of cardiac arrest (Class IIb, LOE B). However, there is insufficient evidence to support the routine use of the LDB in the treatment of cardiac arrest.

Extracorporeal Techniques and Invasive Perfusion Devices

Extracorporeal CPR

For the purpose of these Guidelines, extracorporeal membrane oxygenation (ECMO) and cardiopulmonary bypass are considered together as different forms of extracorporeal CPR (ECPR; an alternative term may be extracorporeal life support or ECLS) when either is used for resuscitation for cardiac arrest. Both are sophisticated

techniques for circulating blood outside the body with or without extracorporeal oxygenation, with the goal of supporting the body's circulation in the absence of an adequately functioning cardiac pump. The initiation of ECPR and the management of a patient on ECPR require highly trained personnel and specialized equipment.

Although there are no data from randomized studies to support the routine use of ECPR, in case series and observational studies the use of ECPR for in-hospital^{107,108} and out-of-hospital^{109–111} cardiac arrest has been associated with improved survival when compared with conventional CPR in patients <75 years old with potentially correctable conditions. However, supportive studies consisted of small numbers of patients, and some had unbalanced comparison groups with respect to age, witnessed arrest, bystander CPR, and the quality of conventional CPR.

There are no randomized studies that compare ECPR with conventional CPR for patients in cardiac arrest. However, data from several case series have demonstrated the feasibility and safety of ECPR in highly specialized centers.^{108,110,111} Observational studies of adults in both the in-hospital¹⁰⁷ and out-of-hospital¹⁰⁹ settings have demonstrated an association between ECPR use and improved survival when compared with conventional CPR in patients with potentially correctable conditions. These studies had small numbers of patients, and some had unbalanced comparison groups with respect to age, witness status, bystander CPR, and the quality of conventional CPR. Please refer to the Pediatrics section for discussion and specific recommendations related to the pediatric population (See Part 14: "Pediatric Advanced Life Support").

There is insufficient evidence to recommend the routine use of ECPR for patients in cardiac arrest. However, in settings where ECPR is readily available, it may be considered when the time without blood flow is brief and the condition leading to the cardiac arrest is reversible (eg, accidental hypothermia drug intoxication) or amenable to heart transplantation (eg, myocarditis) or revascularization (eg, acute myocardial infarction) (Class IIb, LOE C).

Summary

A variety of CPR techniques and devices may improve hemodynamics or short-term survival when used by well-trained providers in selected patients. All of these techniques and devices have the potential to delay chest compressions and defibrillation. In order to prevent delays and maximize efficiency, initial training, ongoing monitoring, and retraining programs should be offered to providers on a frequent and ongoing basis. To date, no adjunct has consistently been shown to be superior to standard conventional (manual) CPR for out-of-hospital basic life support, and no device other than a defibrillator has consistently improved long-term survival from out-of-hospital cardiac arrest.

Disclosures

Guidelines Part 7: CPR Techniques and Devices: Writing Group Disclosures

Writing Group Member	Employment	Research Grant	Other Research		Ownership Interest	Consultant/Advisory	
			Support	Speakers' Bureau/Honoraria		Board	Other
Diana M. Cave	Legacy Health System, Emanuel Hospital, Emergency Services—RN, MSN; Portland Com. College—Institute for Health Prof.-Faculty/Instructor	None	None	None	None	None	None
Raul Gazmuri	North Chicago VA Medical Center—Section Chief, Critical Care and Professor of Medicine	†Volume-Controlled Manual Ventilation during Resuscitation from Cardiac Arrest. Funded by Dessinier Corporation. Funds come to my institution (Rosalind Franklin University(RFU) Vitamin-C Preserves Myocardial Distensibility during Resuscitation from CA. Funded by Maribor University, Slovenia. Funds come to my institution (RFU)	None	None	†Patent titled "Facilitation of Resuscitation from Cardiac Arrest by Erythropoietin" (pending)	None	None
Charles W. Otto	University of Arizona—Professor	None	None	None	None	None	None
Vinay M. Nadkarni	University of Pennsylvania/The Children's Hospital of Philadelphia—Attending Physician, Departement of Anesthesia, Critical Care and Pediatrics	None	None	None	None	None	*Voluntary (Unpaid) member of Data Safety Monitoring Committee for Automated CPR device trial
Adam Cheng	British Columbia Children's Hospital: University Affiliated—Director, Pediatric Simulation Program	†American Heart Association RFP - educational grant. Money comes to my institution, and is distributed to our group of collaborative pediatric hospitals	None	None	None	None	None
Steven C. Brooks	University of Toronto—Clinician-Scientist	†PI-1. Univ. of Toronto Faculty of Medicine New Staff Grant. 01/07/2009–01/07/2010 A pilot study to explore missed opportunities for public access defibrillation in OHCA and to determine the potential impact of emergency medical dispatchers. Role: PI \$10,000 unrestricted grant administered through the research institute 2. University of Toronto Connaught New Staff Matching Grant 2009–2010. 04/05/2009–03/05/2011 Development of Centres of Excellence to Improve Outcomes after OHCA: A Pilot Study. Role: PI \$23,700 unrestricted grant administered through the research institute 3. Ontario Ministry of Health and Long Term Care and the Sunnybrook Medical Services Alternative Funding Plan Association. 04/22/2009–04/21/2010 2008–2009 Alternative Funding Plan Phase III Innovation Fund Project Funding. Project: "Inventing the Future of Post Cardiac Arrest Care: Collaborative Development of Standardized Patient Care Pathways at Sunnybrook Health Sciences Centre." Role: PI \$100,000 unrestricted grant administered through the research institute Co-Investigator 1. National Institutes of Health Slutsky AS (PI) 01/09/2004–01/09/2009 From Bench to Bedside to Curbside. Clinical Research Consortium to improve Resuscitation. Role: Co-Investigator \$2, 454, 201 US 2. Canadian Institute of Health Research Slutsky AS (PI) 01/04/2005–01/10/2010 Epistry component of the Resuscitation Outcomes Consortium. Role: Co-Investigator \$500,001 3. Laerdal Foundation for Acute Medicine Morrison LJ (PI) 01/12/2007–01/12/2010 Centre Grant Program for knowledge translation projects in post resuscitation care. Role: Co-Investigator \$150,000 4. Heart and Stroke Foundation of Canada. Morrison LJ & Dorian P (Co-PI's) 01/12/2007–01/12/2009 Operating Grant in the Area of Resuscitation and Knowledge Transfer for the Strategies in Post-Arrest Care (SPARC) project. Role: Co-Investigator \$200,000	None	None	None	None	None

(Continued)

Guidelines Part 7: CPR Techniques and Devices: Writing Group Disclosures, *Continued*

Writing Group Member	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Ownership Interest	Consultant/Advisory Board	Other
Mohamud Daya	Oregon Health & Science University: Attending Physician—Associate Professor of Emergency Medicine	†PI Resuscitation Outcomes Consortium - Portland Site, NHLBI, grant is awarded directly to the institution (OHSU)	None	*Lectures at local, regional and national meetings, income is directly to me, last lectures CPR update at the Timberline EMS conference, there was no honorarium but conference paid for my lodging Stroke Update in Corvallis at Samaritan Health, Honorarium fee was 500 dollars Advanced 12 lead ECG diagnostic algorithms, Lecture for Philips Healthcare at EMS today, honorarium for 2 lectures was 1000 dollars	*Stock held in the following health care companies; Johnson and Johnson - 250 shares Amgen - 100 shares Roche - 100 shares	*Philips Health Care - Consultant on 12 lead ECG diagnostic algorithms and resuscitation products, no reimbursement for this activity	†I am an EMS medical director for 2 fire departments and one 911 agency, this is a private contract and the money comes directly to me, this is independent of my employment at OHSU which is at an 80% FTE level, my EMS activities are 20% FTE
Robert M. Sutton	The Children's Hospital of Philadelphia—Critical Care Attending	*Unrestricted Research Grant Support through a Center of Excellence Grant from the Laerdal Found	None	None	None	None	
Richard Branson	University of Cincinnati—Associate Professor	None	†SeQual. Sponsor of laboratory study of the use of oxygen concentrators in conjunction with mechanical ventilators for military and mass casualty scenarios. \$40,000. All monies are paid to the Univ. I have no financial interest in the company and do not receive any personal income	†Cardinal - makers of ICU and home care ventilators. I am paid directly for speaking. Newport Medical makers of ICU and home care ventilators. I am paid directly for speaking. *IKARIA - manufactures and distributes inhaled nitric oxide. I am paid directly	None	*Bayer Pharmaceuticals. Treatment of ventilator associated pneumonia	*Kings Daughters Hospital Ashland KY. Paid directly to me
Mary Fran Hazinski	Vanderbilt University School of Nursing—Professor; American Heart Association—Senior Science Editor †Significant AHA compensation for my editing responsibilities—writing and editing of the 2010 AHA Guidelines for CPR and ECC	None	None	None	None	None	None

This table represents the relationships of writing group members that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all members of the writing group are required to complete and submit. A relationship is considered to be "significant" if (a) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (b) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

†Significant.

References

- Ornato JP, Gonzalez ER, Garnett AR, Levine RL, McClung BK. Effect of cardiopulmonary resuscitation compression rate on end-tidal carbon dioxide concentration and arterial pressure in man. *Crit Care Med*. 1988;16:241–245.
- Swenson RD, Weaver WD, Niskanen RA, Martin J, Dahlberg S. Hemodynamics in humans during conventional and experimental methods of cardiopulmonary resuscitation. *Circulation*. 1988;78:630–639.
- Kern KB, Sanders AB, Raife J, Milander MM, Otto CW, Ewy GA. A study of chest compression rates during cardiopulmonary resuscitation in humans: the importance of rate-directed chest compressions. *Arch Intern Med*. 1992;152:145–149.
- Raman J, Saldanha RF, Branch JM, Esmore DS, Spratt PM, Farnsworth AE, Harrison GA, Chang VP, Shanahan MX. Open cardiac compression in the postoperative cardiac intensive care unit. *Anaesth Intensive Care*. 1989;17:129–135.
- Anthi A, Tzelepis GE, Alivizatos P, Michalis A, Palatianos GM, Geroulanos S. Unexpected cardiac arrest after cardiac surgery: incidence, predisposing causes, and outcome of open chest cardiopulmonary resuscitation. *Chest*. 1998;113:15–19.
- Pottle A, Bullock I, Thomas J, Scott L. Survival to discharge following Open Chest Cardiac Compression (OCCC): a 4-year retrospective audit in a cardiothoracic specialist centre—Royal Brompton and Harefield NHS Trust, United Kingdom. *Resuscitation*. 2002;52:269–272.
- Takino M, Okada Y. The optimum timing of resuscitative thoracotomy for non-traumatic out-of-hospital cardiac arrest. *Resuscitation*. 1993;26:69–74.
- Boczar ME, Howard MA, Rivers EP, Martin GB, Horst HM, Lewandowski C, Tomlanovich MC, Nowak RM. A technique revisited: hemodynamic comparison of closed- and open-chest cardiac massage during human cardiopulmonary resuscitation. *Crit Care Med*. 1995;23:498–503.

9. Hachimi-Idrissi S, Leeman J, Hubloue Y, Huyghens L, Corne L. Open chest cardiopulmonary resuscitation in out-of-hospital cardiac arrest. *Resuscitation*. 1997;35:151–156.
10. Calinas-Correia J, Phair I. Physiological variables during open chest cardiopulmonary resuscitation: results from a small series. *J Accid Emerg Med*. 2000;17:201–204.
11. Fialka C, Sebok C, Kemetzhofer P, Kwasny O, Sterz F, Vecsei V. Open-chest cardiopulmonary resuscitation after cardiac arrest in cases of blunt chest or abdominal trauma: a consecutive series of 38 cases. *J Trauma*. 2004;57:809–814.
12. Powell DW, Moore EE, Cothren CC, Ciesla DJ, Burch JM, Moore JB, Johnson JL. Is emergency department resuscitative thoracotomy futile care for the critically injured patient requiring prehospital cardiopulmonary resuscitation? *J Am Coll Surg*. 2004;199:211–215.
13. Sheppard FR, Cothren CC, Moore EE, Orfanakis A, Ciesla DJ, Johnson JL, Burch JM. Emergency department resuscitative thoracotomy for non-torso injuries. *Surgery*. 2006;139:574–576.
14. Seamon MJ, Fisher CA, Gaughan JP, Kulp H, Dempsey DT, Goldberg AJ. Emergency department thoracotomy: survival of the least expected. *World J Surg*. 2008;32:604–612.
15. Powell RW, Gill EA, Jurkovich GJ, Ramenofsky ML. Resuscitative thoracotomy in children and adolescents. *Am Surg*. 1988;54:188–191.
16. Rothenberg SS, Moore EE, Moore FA, Baxter BT, Moore JB, Cleveland HC. Emergency Department thoracotomy in children—a critical analysis. *J Trauma*. 1989;29:1322–1325.
17. Sack JB, Kesselbrenner MB, Jarrad A. Interposed abdominal compression-cardiopulmonary resuscitation and resuscitation outcome during asystole and electromechanical dissociation. *Circulation*. 1992;86:1692–1700.
18. Sack JB, Kesselbrenner MB, Bregman D. Survival from in-hospital cardiac arrest with interposed abdominal counterpulsation during cardiopulmonary resuscitation. *JAMA*. 1992;267:379–385.
19. Babbs CF. Interposed abdominal compression CPR: a comprehensive evidence based review. *Resuscitation*. 2003;59:71–82.
20. Babbs CF. Simplified meta-analysis of clinical trials in resuscitation. *Resuscitation*. 2003;57:245–255.
21. Mateer JR, Stueven HA, Thompson BM, Aprahamian C, Darin JC. Pre-hospital IAC-CPR versus standard CPR: paramedic resuscitation of cardiac arrests. *Am J Emerg Med*. 1985;3:143–146.
22. Waldman PJ, Walters BL, Grunau CF. Pancreatic injury associated with interposed abdominal compressions in pediatric cardiopulmonary resuscitation. *Am J Emerg Med*. 1984;2:510–512.
23. Miller B, Cohen A, Serio A, Bettock D. Hemodynamics of cough cardiopulmonary resuscitation in a patient with sustained torsades de pointes/ventricular flutter. *J Emerg Med*. 1994;12:627–632.
24. Keeble W, Tymchak WJ. Triggering of the Bezold Jarisch Reflex by reperfusion during primary PCI with maintenance of consciousness by cough CPR: a case report and review of pathophysiology. *J Invasive Cardiol*. 2008;20:E239–E242.
25. Niemann JT, Rosborough J, Hausknecht M, Brown D, Criley JM. Cough-CPR: documentation of systemic perfusion in man and in an experimental model: a “window” to the mechanism of blood flow in external CPR. *Crit Care Med*. 1980;8:141–146.
26. Saba SE, David SW. Sustained consciousness during ventricular fibrillation: case report of cough cardiopulmonary resuscitation. *Cathet Cardiovasc Diagn*. 1996;37:47–48.
27. Criley JM, Blaufuss AH, Kissel GL. Cough-induced cardiac compression: self-administered form of cardiopulmonary resuscitation. *JAMA*. 1976;236:1246–1250.
28. Wei JY, Greene HL, Weisfeldt ML. Cough-facilitated conversion of ventricular tachycardia. *Am J Cardiol*. 1980;45:174–176.
29. Caldwell G, Millar G, Quinn E, Vincent R, Chamberlain DA. Simple mechanical methods for cardioversion: defence of the precordial thump and cough version. *Br Med J (Clin Res Ed)*. 1985;291:627–630.
30. Miller B, Lesnefsky E, Heyborne T, Schmidt B, Freeman K, Breckinridge S, Kelley K, Mann D, Reiter M. Cough-cardiopulmonary resuscitation in the cardiac catheterization laboratory: hemodynamics during an episode of prolonged hypotensive ventricular tachycardia. *Cathet Cardiovasc Diagn*. 1989;18:168–171.
31. Rieser MJ. The use of cough-CPR in patients with acute myocardial infarction. *J Emerg Med*. 1992;10:291–293.
32. Petelenz T, Iwinski J, Chlebowczyk J, Czyz Z, Flak Z, Fiutowski L, Zaorski K, Zeman S. Self-administered cough cardiopulmonary resuscitation (c-CPR) in patients threatened by MAS events of cardiovascular origin. *Wiad Lek*. 1998;51(7–8):326–336.
33. Girsky MJ, Criley JM. Images in cardiovascular medicine. Cough cardiopulmonary resuscitation revisited. *Circulation*. 2006;114:e530–531.
34. Mazer SP, Weisfeldt M, Bai D, Cardinale C, Arora R, Ma C, Sciacca RR, Chong D, Rabbani LE. Reverse CPR: a pilot study of CPR in the prone position. *Resuscitation*. 2003;57:279–285.
35. Sun WZ, Huang FY, Kung KL, Fan SZ, Chen TL. Successful cardiopulmonary resuscitation of two patients in the prone position using reversed precordial compression. *Anesthesiology*. 1992;77:202–204.
36. Tobias JD, Mencio GA, Atwood R, Gurwitz GS. Intraoperative cardiopulmonary resuscitation in the prone position. *J Pediatr Surg*. 1994;29:1537–1538.
37. Brown J, Rogers J, Soar J. Cardiac arrest during surgery and ventilation in the prone position: a case report and systematic review. *Resuscitation*. 2001;50:233–238.
38. Shuster M, Lim SH, Deakin CD, Kleinman ME, Koster RW, Morrison LJ, Nolan JP, Sayre MR; on behalf of the CPR Techniques and Devices Collaborators. Part 7: CPR techniques and devices: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Circulation*. 2010;122(suppl 2):S338–S344.
39. Pellis T, Kette F, Lovisa D, Franceschino E, Magagnin L, Mercante WP, Kohl P. Utility of pre-cordial thump for treatment of out of hospital cardiac arrest: a prospective study. *Resuscitation*. 2009;80:17–23.
40. Bornemann C, Scherf D. Electrocardiogram of the month. Paroxysmal ventricular tachycardia abolished by a blow to the precordium. *Dis Chest*. 1969;56:83–84.
41. Dale KM, Lertsburapa K, Kluger J, White CM. Moxifloxacin and torsade de pointes. *Ann Pharmacother*. 2007;41:336–340.
42. De Maio VJ, Stiell IG, Spaite DW, Ward RE, Lyver MB, Field BJ III, Munkley DP, Wells GA. CPR-only survivors of out-of-hospital cardiac arrest: implications for out-of-hospital care and cardiac arrest research methodology. *Ann Emerg Med*. 2001;37:602–608.
43. Pennington JE, Taylor J, Lown B. Chest thump for reverting ventricular tachycardia. *N Engl J Med*. 1970;283:1192–1195.
44. Rahner E, Zeh E. Die Regularisierung von Kammertachykardien durch präkordialen Faustschlag. (“The Regularization of Ventricular Tachycardias by Precordial Thumping.”) *Med Welt*. 1978;29:1659–1663.
45. Amir O, Schliamser JE, Nemer S, Arie M. Ineffectiveness of precordial thump for cardioversion of malignant ventricular tachyarrhythmias. *Pacing Clin Electrophysiol*. 2007;30:153–156.
46. Haman L, Parizek P, Vojacek J. Precordial thump efficacy in termination of induced ventricular arrhythmias. *Resuscitation*. 2009;80:14–16.
47. Ahmar W, Morley P, Marasco S, Chan W, Aggarwal A. Sternal fracture and osteomyelitis: an unusual complication of a precordial thump. *Resuscitation*. 2007;75:540–542.
48. Miller J, Tresch D, Horwitz L, Thompson BM, Aprahamian C, Darin JC. The precordial thump. *Ann Emerg Med*. 1984;13(9 Pt 2):791–794.
49. Muller GI, Ulmer HE, Bauer JA. Complications of chest thump for termination of supraventricular tachycardia in children. *Eur J Pediatr*. 1992;151:12–14.
50. Chan L, Reid C, Taylor B. Effect of three emergency pacing modalities on cardiac output in cardiac arrest due to ventricular asystole. *Resuscitation*. 2002;52:117–119.
51. Dowdle JR. Ventricular standstill and cardiac percussion. *Resuscitation*. 1996;32:31–32.
52. Eich C, Bleckmann A, Schwarz SK. Percussion pacing—an almost forgotten procedure for haemodynamically unstable bradycardias? A report of three case studies and review of the literature. *Br J Anaesth*. 2007;98:429–433.
53. Eich C, Bleckmann A, Paul T. Percussion pacing in a three-year-old girl with complete heart block during cardiac catheterization. *Br J Anaesth*. 2005;95:465–467.
54. Iseri LT, Allen BJ, Baron K, Brodsky MA. Fist pacing, a forgotten procedure in bradysystolic cardiac arrest. *Am Heart J*. 1987;113:1545–1550.
55. Tucker KJ, Shaburhivili TS, Gedeveanishvili AT. Manual external (fist) pacing during high-degree atrioventricular block: a lifesaving intervention. *Am J Emerg Med*. 1995;13:53–54.
56. Zeh E, Rahner E. [The manual extrathoracic stimulation of the heart. Technique and effect of the precordial thump (author’s transl)]. *Z Kardiol*. 1978;67:299–304.

57. Weiss SJ, Ernst AA, Jones R, Ong M, Filbrun T, Augustin C, Barnum M, Nick TG. Automatic transport ventilator versus bag valve in the EMS setting: a prospective, randomized trial. *South Med J*. 2005;98:970–976.
58. Johannigman JA, Branson RD, Johnson DJ, Davis K Jr, Hurst JM. Out-of-hospital ventilation: bag-valve device vs transport ventilator. *Acad Emerg Med*. 1995;2:719–724.
59. Noordergraaf GJ, van Dun PJ, Kramer BP, Schors MP, Hornman HP, de Jong W, Noordergraaf A. Can first responders achieve and maintain normocapnia when sequentially ventilating with a bag-valve device and two oxygen-driven resuscitators? A controlled clinical trial in 104 patients. *Eur J Anaesthesiol*. 2004;21:367–372.
60. Hevesi ZG, Thrush DN, Downs JB, Smith RA. Cardiopulmonary resuscitation: effect of CPAP on gas exchange during chest compressions. *Anesthesiology*. 1999;90:1078–1083.
61. Cohen TJ, Goldner BG, Maccaro PC, Ardito AP, Trazzera S, Cohen MB, Dibs SR. A comparison of active compression-decompression cardiopulmonary resuscitation with standard cardiopulmonary resuscitation for cardiac arrests occurring in the hospital. *N Engl J Med*. 1993;329:1918–1921.
62. Lurie KG, Shultz JJ, Callahan ML, Schwab TM, Gisch T, Rector T, Frascone RJ, Long L. Evaluation of active compression-decompression CPR in victims of out-of-hospital cardiac arrest. *JAMA*. 1994;271:1405–1411.
63. Tucker KJ, Galli F, Savitt MA, Kahsai D, Bresnahan L, Redberg RF. Active compression-decompression resuscitation: effect on resuscitation success after in-hospital cardiac arrest. *J Am Coll Cardiol*. 1994;24:201–209.
64. Plaisance P, Adnet F, Vicaut E, Hennequin B, Magne P, Prudhomme C, Lambert Y, Cantineau JP, Leopold C, Ferracci C, Gizzi M, Payen D. Benefit of active compression-decompression cardiopulmonary resuscitation as a prehospital advanced cardiac life support: a randomized multicenter study. *Circulation*. 1997;95:955–961.
65. Plaisance P, Lurie KG, Vicaut E, Adnet F, Petit JL, Epain D, Ecollan P, Gruat R, Cavagna P, Biens J, Payen D. A comparison of standard cardiopulmonary resuscitation and active compression-decompression resuscitation for out-of-hospital cardiac arrest. French Active Compression-Decompression Cardiopulmonary Resuscitation Study Group. *N Engl J Med*. 1999;341:569–575.
66. He Q, Wan Z, Wang L. [Random control trial of the efficacy of cardiopump on pre-hospital cardiac arrest]. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue*. 2003;15:292–294.
67. Lafuente-Lafuente C, Melero-Bascones M. Active chest compression-decompression for cardiopulmonary resuscitation. *Cochrane Database Syst Rev*. 2004;:CD002751.
68. Mauer D, Schneider T, Dick W, Wilhelm A, Elich D, Mauer M. Active compression-decompression resuscitation: a prospective, randomized study in a two-tiered EMS system with physicians in the field. *Resuscitation*. 1996;33:125–134.
69. Stiell IG, Hebert PC, Wells GA, Laupacis A, Vandemheen K, Dreyer JF, Eisenhauer MA, Gibson J, Higginson LA, Kirby AS, Mahon JL, Maloney JP, Weitzman BN. The Ontario trial of active compression-decompression cardiopulmonary resuscitation for in-hospital and pre-hospital cardiac arrest. *JAMA*. 1996;275:1417–1423.
70. Goralski M, Villegier JL, Cami G, Linassier P, Guilles-Des-Buttes P, Fabbri P, Venot P, Tazarourte K, Cami M. Evaluation of active compression-decompression cardiopulmonary resuscitation in out-of-hospital cardiac arrest. *Resuscitation*. 1998;7:543–550.
71. Skogvoll E, Wik L. Active compression-decompression cardiopulmonary resuscitation: a population-based, prospective randomised clinical trial in out-of-hospital cardiac arrest. *Resuscitation*. 1999;42:163–172.
72. Schwab TM, Callahan ML, Madsen CD, Utecht TA. A randomized clinical trial of active compression-decompression CPR vs standard CPR in out-of-hospital cardiac arrest in two cities. *JAMA*. 1995;273:1261–1268.
73. Luiz T, Ellinger K, Denz C. Active compression-decompression cardiopulmonary resuscitation does not improve survival in patients with prehospital cardiac arrest in a physician-manned emergency medical system. *J Cardiothorac Vasc Anesth*. 1996;10:178–186.
74. Nolan J, Smith G, Evans R, McCusker K, Lubas P, Parr M, Baskett P. The United Kingdom pre-hospital study of active compression-decompression resuscitation. *Resuscitation*. 1998;37:119–125.
75. Arntz HR, Agrawal R, Richter H, Schmidt S, Rescheleit T, Menges M, Burbach H, Schroder J, Schultheiss HP. Phased chest and abdominal compression-decompression versus conventional cardiopulmonary resuscitation in out-of-hospital cardiac arrest. *Circulation*. 2001;104:768–772.
76. Plaisance P, Lurie KG, Payen D. Inspiratory impedance during active compression-decompression cardiopulmonary resuscitation: a randomized evaluation in patients in cardiac arrest. *Circulation*. 2000;101:989–994.
77. Plaisance P, Soleil C, Lurie KG, Vicaut E, Ducros L, Payen D. Use of an inspiratory impedance threshold device on a facemask and endotracheal tube to reduce intrathoracic pressures during the decompression phase of active compression-decompression cardiopulmonary resuscitation. *Crit Care Med*. 2005;33:990–994.
78. Wolcke BB, Mauer DK, Schoefmann MF, Teichmann H, Provo TA, Lindner KH, Dick WF, Aeppli D, Lurie KG. Comparison of standard cardiopulmonary resuscitation versus the combination of active compression-decompression cardiopulmonary resuscitation and an inspiratory impedance threshold device for out-of-hospital cardiac arrest. *Circulation*. 2003;108:2201–2205.
79. Plaisance P, Lurie KG, Vicaut E, Martin D, Gueugniaud PY, Petit JL, Payen D. Evaluation of an impedance threshold device in patients receiving active compression-decompression cardiopulmonary resuscitation for out of hospital cardiac arrest. *Resuscitation*. 2004;61:265–271.
80. Aufderheide TP, Pirralo RG, Provo TA, Lurie KG. Clinical evaluation of an inspiratory impedance threshold device during standard cardiopulmonary resuscitation in patients with out-of-hospital cardiac arrest. *Crit Care Med*. 2005;33:734–740.
81. Pirralo RG, Aufderheide TP, Provo TA, Lurie KG. Effect of an inspiratory impedance threshold device on hemodynamics during conventional manual cardiopulmonary resuscitation. *Resuscitation*. 2005;66:13–20.
82. Thayne RC, Thomas DC, Neville JD, Van Dellen A. Use of an impedance threshold device improves short-term outcomes following out-of-hospital cardiac arrest. *Resuscitation*. 2005;67:103–108.
83. Cabrini L, Beccaria P, Landoni G, Biondi-Zoccai GG, Sheiban I, Cristofolini M, Fochi O, Maj G, Zangrillo A. Impact of impedance threshold devices on cardiopulmonary resuscitation: a systematic review and meta-analysis of randomized controlled studies. *Crit Care Med*. 2008;36:1625–1632.
84. Aufderheide T, Alexander C, Lick C, Myers B, Romig L, Vartanian L, Stothert J, McKnite S, Matsuura T, Yannopoulos D, Lurie K. From laboratory science to six emergency medical services systems: new understanding of the physiology of cardiopulmonary resuscitation increase survival rates after cardiac arrest. *Crit Care Med*. 2008;36(11[Suppl.]):S397–S404.
85. Aufderheide TP, Yannopoulos D, Lick CJ, Myers B, Romig LA, Stothert JC, Barnard J, Vartanian L, Pilgrim AJ, Benditt DG. Implementing the 2005 American Heart Association Guidelines improves outcomes after out-of-hospital cardiac arrest. April 26, 2010. doi:10.1016/j.hrthm.2010.04.022. Available at: <http://www.heartrhythmjournal.com/article/PIIS1547527110003978/fulltext>.
86. Hinchey PR, Myers JB, Lewis R, De Maio VJ, Reyer E, Licatose D, Zalkin J, Snyder G. Improved out-of-hospital cardiac arrest survival after the sequential implementation of 2005 AHA guidelines for compressions, ventilations, and induced hypothermia: the Wake County experience. *Ann Emerg Med*. 2010. April 1, 2010. doi:10.1016/j.annemergmed.2010.01.036. Available at: [http://www.annemergmed.com/article/S0196-0644\(10\)00116-2/fulltext](http://www.annemergmed.com/article/S0196-0644(10)00116-2/fulltext).
87. Dickinson ET, Verdile VP, Schneider RM, Salluzzo RF. Effectiveness of mechanical versus manual chest compressions in out-of-hospital cardiac arrest resuscitation: a pilot study. *Am J Emerg Med*. 1998;16:289–292.
88. McDonald JL. Systolic and mean arterial pressures during manual and mechanical CPR in humans. *Ann Emerg Med*. 1982;11:292–295.
89. Ward KR, Menegazzi JJ, Zelenak RR, Sullivan RJ, McSwain NE Jr. A comparison of chest compressions between mechanical and manual CPR by monitoring end-tidal PCO2 during human cardiac arrest. *Ann Emerg Med*. 1993;22:669–674.
90. Taylor GJ, Rubin R, Tucker M, Greene HL, Rudikoff MT, Weisfeldt ML. External cardiac compression: a randomized comparison of mechanical and manual techniques. *JAMA*. 1978;240:644–646.
91. Wang HC, Chiang WC, Chen SY, Ke YL, Chi CL, Yang CW, Lin PC, Ko PC, Wang YC, Tsai TC, Huang CH, Hsiung KH, Ma MH, Chen SC, Chen WJ, Lin FY. Video-recording and time-motion analyses of manual versus mechanical cardiopulmonary resuscitation during ambulance transport. *Resuscitation*. 2007;74:453–460.

92. Axelsson C, Nestin J, Svensson L, Axelsson AB, Herlitz J. Clinical consequences of the introduction of mechanical chest compression in the EMS system for treatment of out-of-hospital cardiac arrest—a pilot study. *Resuscitation*. 2006;71:47–55.
93. Steen S, Liao Q, Pierre L, Paskevicius A, Sjoberg T. Evaluation of LUCAS, a new device for automatic mechanical compression and active decompression resuscitation. *Resuscitation*. 2002;55:285–299.
94. Steen S, Sjoberg T, Olsson P, Young M. Treatment of out-of-hospital cardiac arrest with LUCAS, a new device for automatic mechanical compression and active decompression resuscitation. *Resuscitation*. 2005;67:25–30.
95. Larsen AI, Hjernevik AS, Ellingsen CL, Nilsen DW. Cardiac arrest with continuous mechanical chest compression during percutaneous coronary intervention: a report on the use of the LUCAS device. *Resuscitation*. 2007;75:454–459.
96. Deakin CD, O'Neill JF, Tabor T. Does compression-only cardiopulmonary resuscitation generate adequate passive ventilation during cardiac arrest? *Resuscitation*. 2007;75:53–59.
97. Bonnemeier H, Olivecrona G, Simonis G, Gotberg M, Weitz G, Iblher P, Gerling I, Schunkert H. Automated continuous chest compression for in-hospital cardiopulmonary resuscitation of patients with pulseless electrical activity: a report of five cases. *Int J Cardiol*. 2009;136:e39–e50.
98. Wagner H, Terkelsen CJ, Friberg H, Harnek J, Kern K, Lassen JF, Olivecrona GK. Cardiac arrest in the catheterisation laboratory: a 5-year experience of using mechanical chest compressions to facilitate PCI during prolonged resuscitation efforts. *Resuscitation*. 2009.
99. Wirth S, Korner M, Treitl M, Linsenmaier U, Leidel BA, Jaschkowitz T, Reiser MF, Kanz KG. Computed tomography during cardiopulmonary resuscitation using automated chest compression devices: an initial study. *Eur Radiol*. 2009;19:1857–1866.
100. Timmerman S, Cardoso LF, Ramires JA, Halperin H. Improved hemodynamic performance with a novel chest compression device during treatment of in-hospital cardiac arrest. *Resuscitation*. 2004;61:273–280.
101. Casner M, Andersen D, Isaacs SM. The impact of a new CPR assist device on rate of return of spontaneous circulation in out-of-hospital cardiac arrest. *Prehosp Emerg Care*. 2005;9:61–67.
102. Ong ME, Ornato JP, Edwards DP, Dhindsa HS, Best AM, Ines CS, Hickey S, Clark B, Williams DC, Powell RG, Overton JL, Peberdy MA. Use of an automated, load-distributing band chest compression device for out-of-hospital cardiac arrest resuscitation. *JAMA*. 2006;295:2629–2637.
103. Steinmetz J, Barnung S, Nielsen SL, Risom M, Rasmussen LS. Improved survival after an out-of-hospital cardiac arrest using new guidelines. *Acta Anaesthesiol Scand*. 2008;52:908–913.
104. Hallstrom A, Rea TD, Sayre MR, Christenson J, Anton AR, Mosesso VN Jr, Van Ottingham L, Olsufka M, Pennington S, White LJ, Yahn S, Husar J, Morris MF, Cobb LA. Manual chest compression vs use of an automated chest compression device during resuscitation following out-of-hospital cardiac arrest: a randomized trial. *JAMA*. 2006;295:2620–2628.
- 104A. Hallstrom A, Rea TD, Sayre MR, Christenson J, Cobb LA, Mosesso VN Jr, Anton AR. The ASPIRE trial investigators respond to inhomogeneity and temporal effects assertion. *Am J Emerg Med*. August 16, 2010. doi:10.1016/j.ajem.2010.07.001. Available at: [http://www.ajem-journal.com/article/S0735-6757\(10\)00307-4/fulltext](http://www.ajem-journal.com/article/S0735-6757(10)00307-4/fulltext).
105. Paradis N, Young G, Lemeshow S, Brewer J, Halperin H. Inhomogeneity and temporal effects in ASPIRE - An Exception from Consent Trial Terminated Early. *Am J Emerg Med*. In Press.
106. Tomte O, Sunde K, Lorentz T, Auestad B, Souders C, Jensen J, Wik L. Advanced life support performance with manual and mechanical chest compressions in a randomized, multicentre manikin study. *Resuscitation*. 2009;80:1152–1157.
107. Chen YS, Lin JW, Yu HY, Ko WJ, Jerng JS, Chang WT, Chen WJ, Huang SC, Chi NH, Wang CH, Chen LC, Tsai PR, Wang SS, Hwang JJ, Lin FY. Cardiopulmonary resuscitation with assisted extracorporeal life-support versus conventional cardiopulmonary resuscitation in adults with in-hospital cardiac arrest: an observational study and propensity analysis. *Lancet*. 2008;372:554–561.
108. Athanasuleas CL, Buckberg GD, Allen BS, Beyersdorf F, Kirsh MM. Sudden cardiac death: directing the scope of resuscitation towards the heart and brain. *Resuscitation*. 2006;70:44–51.
109. Tanno K, Itoh Y, Takeyama Y, Nara S, Mori K, Asai Y. Utstein style study of cardiopulmonary bypass after cardiac arrest. *Am J Emerg Med*. 2008;26:649–654.
110. Chen YS, Yu HY, Huang SC, Lin JW, Chi NH, Wang CH, Wang SS, Lin FY, Ko WJ. Extracorporeal membrane oxygenation support can extend the duration of cardiopulmonary resuscitation. *Crit Care Med*. 2008;36:2529–2535.
111. Nagao K, Kikushima K, Watanabe K, Tachibana E, Tominaga Y, Tada K, Ishii M, Chiba N, Kasai A, Soga T, Matsuzaki M, Nishikawa K, Tateda Y, Ikeda H, Yagi T. Early induction of hypothermia during cardiac arrest improves neurological outcomes in patients with out-of-hospital cardiac arrest who undergo emergency cardiopulmonary bypass and percutaneous coronary intervention. *Circ J*. 2010;74:77–85.

KEY WORDS: cardiac arrest ■ cardiopulmonary resuscitation ■ emergency ■ ventricular fibrillation

**Part 7: CPR Techniques and Devices: 2010 American Heart Association Guidelines for
Cardiopulmonary Resuscitation and Emergency Cardiovascular Care**

Diana M. Cave, Raul J. Gazmuri, Charles W. Otto, Vinay M. Nadkarni, Adam Cheng, Steven C. Brooks, Mohamud Daya, Robert M. Sutton, Richard Branson and Mary Fran Hazinski

Circulation. 2010;122:S720-S728

doi: 10.1161/CIRCULATIONAHA.110.970970

Circulation is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2010 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

http://circ.ahajournals.org/content/122/18_suppl_3/S720

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation* is online at:
<http://circ.ahajournals.org/subscriptions/>