

Out-of-Hospital Cardiac Arrest

An Underlying Reversible Cause

ECG CHALLENGE

An 84-year-old man called the emergency service from his home because of recent chest pain associated with dyspnea. He had a long-standing history of coronary artery disease and chronic obstructive pulmonary disease. One minute after the basic life support team arrived, the patient went into cardiac arrest. The rescuers deployed their automated external defibrillator, and 1 electric shock was delivered. Thereafter, the patient received cardiopulmonary resuscitation. The prehospital emergency physician arrived at the scene 10 minutes later. At this time, the patient had pulseless electric activity (PEA). The ECG monitoring (lead 2) during cardiopulmonary resuscitation is shown in Figure 1. He received 1 mg adrenaline, continuous chest compressions, and orotracheal intubation. A few minutes later, the patient had recovered spontaneous circulation. Cardiac and pulmonary auscultations were normal. The patient was connected to a ventilator, which displayed normal pressure levels. He was then transported to the emergency department without any new clinical complications.

What abnormalities are present in the ECG, and might they relate to a particular cause of the cardiac arrest?

Jimmy Robert, MD
Clément Derkenne, MD
Daniel Jost, MD
Jean-Pierre Tourtier, PhD




Figure 1. The 2 tracings represent the ECG and the thoracic impedance variation. The hands-off times reveal the presence of electric alternans of the ECG, which persisted during the chest compressions.

Please turn the page to read the diagnosis.

Correspondence to: Clément Derkenne, MD, Paris Fire Brigade, Emergency Medical Department, 1, Place Jules Renard, 75017 Paris, France. E-mail clement.derkenne@pompierparis.fr

Key Words: electrocardiography ■ heart arrest ■ pneumothorax

© 2017 American Heart Association, Inc.

RESPONSE TO ECG CHALLENGE

The ECG in Figure 1 shows 2 tracings; lead 2 of the cardiac ECG and the thoracic impedance variation display. The patient's thoracic impedance was measured with a continuous low electric current generated by the monitor via the defibrillation electrodes (LifePack15, Physio-Control, Seattle, WA). Its variation represents chest movement, especially when the patient receives chest compressions. Lead 2 can be best interpreted during periods without chest compressions (those with a flat impedance curve). The first hands-off time reveals PEA with a rhythm that is not strictly regular, without a visible P wave, and with a frequency of 60 bpm, a QRS interval of 150 milliseconds, and a QTc interval of 300 milliseconds. The most important observation is the presence of electric alternans. Electric alternans is a rarely observed cardiac phenomenon in which the ECG tracing shows alternation of the QRS complex amplitude, morphology, or axis, most often with a 1-in-2 periodicity. Although this electric phenomenon is not pathognomonic, it is usually attributable to pericardial effusion, hemothorax, or pneumothorax.¹

Several mechanisms have been proposed to explain the phenomenon of electric alternans, including swinging of the heart, which causes the electric axis to change (as in pericardial effusion), and changes to the volume of the thoracic cavity during respiratory movements. At the hospital, the patient received a thoracic computed tomography scan that showed complete and compressive right pneumothorax (Figure 2). Exsufflation was immediately performed, followed by chest tube insertion, whereupon the electric alternans disappeared immediately. On the basis of the history of atraumatic emphysema, the rupture of an emphysema bleb was believed to be the most likely explanation for the pneumothorax. Three days later, the patient was discharged from the intensive care unit.

This is the first report of electric alternans observed during resuscitation for an out-of-hospital cardiac arrest. The current guidelines for advanced life support in patients with out-of-hospital cardiac arrest recommend 2-minute cardiopulmonary resuscitation cycles, which are usually punctuated by analysis of the rhythm and classification as shockable or nonshockable. PEA is the most common nonshockable rhythm associated with a correctable cause.² PEA can be associated with several different causes, each which may have a specific manifestation on the ECG such as electric alternans in cases of tamponade and pneumothorax, Osborn (J) wave in hypothermia, or widening QRS in tricyclic antidepressant intoxication.³ Each specific type of PEA requires a specific treatment.³

In this case, if electric alternans had been recognized earlier by the emergency physician, the patient could have received a prehospital pulmonary and cardiac ultrasound examination and potentially could have been diagnosed and treated promptly in the field. Of note, there was neither ipsilateral nor contralateral decreased air entry on auscultation. The normality of auscultation was unexpected, but the literature shows normal auscultation in up to 42% of cases of pneumothorax.

In summary, this is a case of electric alternans observed during resuscitation for out-of-hospital cardiac arrest in a subject with a history of emphysema. The recognition of a correctable cause on ECG has the potential of improving prognosis in some patients who experience out-of-hospital cardiac arrest.

ACKNOWLEDGMENTS

The authors thank Fabrice Metayer for his help with the Physio-Control aspects of this article.



Figure 2. Computed tomography scan revealing a complete right pneumothorax (asterisk).

DISCLOSURES

None.

AFFILIATIONS

From Paris Fire Brigade, Emergency Medical Department, France (J.R., C.D., D.J., J.-P.T.); and Sudden Death Expertise Center, INSERM U.970, Paris, France (D.J.).

FOOTNOTES

Circulation is available at <http://circ.ahajournals.org>.

REFERENCES

1. Surawicz B, Fisch C. Cardiac alternans: diverse mechanisms and clinical manifestations. *J Am Coll Cardiol*. 1992;20:483–499.
2. Myerburg RJ, Halperin H, Egan DA, Boineau R, Chugh SS, Gillis AM, Goldhaber JL, Lathrop DA, Liu P, Niemann JT, Ornato JP, Sopko G, Van Eyk JE, Walcott GP, Weisfeldt ML, Wright JD, Zipes DP. Pulseless electric activity: definition, causes, mechanisms, management, and research priorities for the next decade: report from a National Heart, Lung, and Blood Institute workshop. *Circulation*. 2013;128:2532–2541. doi: 10.1161/CIRCULATIONAHA.113.004490.
3. Jentzer JC, Clements CM, Wright RS, White RD, Jaffe AS. Improving survival from cardiac arrest: a review of contemporary practice and challenges. *Ann Emerg Med*. 2016;68:678–689. doi: 10.1016/j.annemergmed.2016.05.022.

Out-of-Hospital Cardiac Arrest: An Underlying Reversible Cause
Jimmy Robert, Clément Derkenne, Daniel Jost and Jean-Pierre Tourtier

Circulation. 2017;135:2564-2566

doi: 10.1161/CIRCULATIONAHA.117.028104

Circulation is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2017 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circ.ahajournals.org/content/135/25/2564>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation* is online at:
<http://circ.ahajournals.org/subscriptions/>