

Use of Sildenafil (Viagra) in Patients With Cardiovascular Disease

Writing Group Members

Melvin D. Cheitlin, MD, FACC, Cochair; Adolph M. Hutter, Jr, MD, MACC, Cochair;
Ralph G. Brindis, MD, MPH, FACC; Peter Ganz, MD, FACC; Sanjay Kaul, MD;
Richard O. Russell, Jr, MD, FACC; Randall M. Zusman, MD, FACC*

Technology and Practice Executive Committee

James S. Forrester, MD, FACC, Chair; Pamela S. Douglas, MD, FACC; David P. Faxon, MD, FACC;
John D. Fisher, MD, FACC; Raymond J. Gibbons, MD, FACC; Jonathan L. Halperin, MD, FACC;
Adolph M. Hutter, Jr, MD, MACC; Judith S. Hochman, MD, FACC; Sanjiv Kaul, MD, FACC*;
William S. Weintraub, MD, FACC; William L. Winters, Jr, MD, MACC; Michael J. Wolk, MD, FACC

Executive Summary

The pharmaceutical preparation sildenafil citrate (Viagra) is being widely prescribed as a treatment for male erectile dysfunction, a common problem that in the United States affects between 10 and 30 million men. The introduction of sildenafil has been a valuable contribution to the treatment of erectile dysfunction, which is a relatively common occurrence in patients with cardiovascular disease. This article is written to appropriately caution and not to unduly alarm physicians in their use of sildenafil in patients with heart disease.

Reported cardiovascular side effects in the normal healthy population are typically minor and associated with vasodilatation (ie, headache, flushing, and small decreases in systolic and diastolic blood pressures). However, although their incidence is small, serious cardiovascular events, including significant hypotension, can occur in certain populations at risk. Most at risk are individuals who are concurrently taking organic nitrates. Organic nitrate preparations are commonly prescribed to manage the symptoms of angina pectoris. The coadministration of nitrates and Viagra significantly increases the risk of potentially life-threatening hypotension. Therefore, Viagra should not be prescribed to patients receiving any form of nitrate therapy.

Although definitive evidence is currently lacking, it is possible that a precipitous reduction in blood pressure with nitrate use may occur over the initial 24 hours after a dose of

Viagra. Thus, for patients who experience an acute cardiac ischemic event and who have taken Viagra within the past 24 hours, administration of nitrates should be avoided. In the event that nitrates are given, especially within this critical time interval, it is essential to have the capability to support the patient with fluid resuscitation and α -adrenergic agonists if needed. In patients with recurring angina after Viagra use, other nonnitrate antianginal agents, such as β -blockers, should be considered.

Other patients in whom the use of Viagra is potentially hazardous include those with active coronary ischemia; those with congestive heart failure and borderline low blood volume and low blood pressure status; those with complicated, multidrug, antihypertensive therapy regimens; and those taking medications that may affect the metabolic clearance of Viagra. With respect to patients following complicated multidrug, antihypertensive programs, the randomized studies included a large number of hypertensive patients. However, most patients were controlled with 1 antihypertensive agent, and only a small number were controlled with 3 antihypertensive agents. Until adequate studies are done in these subgroups of patients, sildenafil should be prescribed with caution.

Viagra acts as a selective inhibitor of cyclic GMP (cGMP)-specific phosphodiesterase type 5, resulting in smooth muscle relaxation, vasodilatation, and enhanced penile erection. Although the cardiovascular effects of sildenafil

The ACC/AHA Expert Consensus Document "Use of Sildenafil (Viagra) in Patients With Cardiovascular Disease" was approved by the Board of Trustees of the American College of Cardiology in September 1998 and the American Heart Association Science Advisory and Coordinating Committee in September 1998. A single reprint is available by calling 800-242-8721 (US only) or writing the American Heart Association, Public Information, 7272 Greenville Ave, Dallas, TX 75231-4596. Ask for reprint No. 71-0156. To purchase additional reprints: up to 999 copies, call 800-611-6083 (US only) or fax 413-665-2671; 1000 or more copies, call 214-706-1466, fax 214-691-6342, or E-mail pubauth@heart.org. To make photocopies for personal or educational use, call the Copyright Clearance Center at 978-750-8400.

*Those authors designated with an asterisk have indicated a potential conflict of interest with respect to the topic of this document. They have excused themselves from discussions or the preparation of the text whence this potential conflict would apply.

(*Circulation*. 1999;99:168-177.)

© 1999 American College of Cardiology and American Heart Association, Inc.

Circulation is available at <http://www.circulationaha.org>

Summary Table of Clinical Recommendations

-
- A. Use of Viagra clearly contraindicated
1. Concurrent use of nitrates (see Appendix A)
- B. Cardiovascular effects of Viagra may be potentially hazardous (use dependent on individual clinical assessment)
1. Patients with active coronary ischemia who are not taking nitrates (eg, positive exercise test for ischemia)
 2. Patients with congestive heart failure and borderline low blood pressure and borderline low volume status
 3. Patients on a complicated, multidrug, antihypertensive program
 4. Patients taking drugs that can prolong the half-life of Viagra (see Appendix B)
-

reported in available randomized, controlled clinical trials were relatively minor, heart disease patients represented only a small fraction of studied patients, and patients with heart failure, patients with myocardial infarction or stroke within 6 months, or patients with uncontrolled hypertension were not included in these studies. Thus, there are possible problems in the use of Viagra in these patients that have not been adequately studied.

Given the increasing reports of deaths in which the use of Viagra may be implicated, clinicians need to exercise caution when advising their patients with heart disease about taking this medication. Specific recommendations regarding sildenafil (Viagra) and the cardiac patient are summarized in the following Table.

I. Preamble

The present document is an expert consensus. This type of document is intended to inform practitioners, payers, and other interested parties of the opinion of the American College of Cardiology (ACC) concerning evolving areas of clinical practice and/or technologies that are widely available or are new to the practice community. Topics chosen for coverage by Expert Consensus Documents are so designated because the evidence base and experience with the technology or clinical practice are not sufficiently well developed to be evaluated by the formal ACC/American Heart Association (AHA) Practice Guidelines process. Thus, the reader should view the Expert Consensus Documents as the best attempt of the ACC to inform and guide clinical practice in areas in which rigorous evidence is not yet available. Where feasible, Expert Consensus Documents will include indications and contraindications. Some topics covered by Expert Consensus Documents will be addressed subsequently by the ACC/AHA Practice Guideline process.

A. Sildenafil (Viagra) Use for Erectile Dysfunction

Male erectile dysfunction defined as "the inability to attain and/or maintain penile erection sufficient for satisfactory sexual performance"¹ is a common problem in the United States affecting between 10 and 30 million men.^{2,3} Sexual dysfunction in men after the diagnosis of coronary artery disease or a myocardial infarction is common. Most is due to fear that the exertion of sexual activity will precipitate another myocardial infarction, but 10% to 15% is due to organic causes of impotence.⁴ Approximately 5.5 million

men take nitrates on a regular basis for angina pectoris,⁵ and another half a million will experience a heart attack annually and are potential candidates for nitrate therapy.⁶ Sildenafil is potentially contraindicated in as many as 6 million patients.

The introduction of sildenafil citrate (Viagra), a drug that acts as a selective inhibitor of cGMP-specific phosphodiesterase type 5 (PDE5), which results in smooth muscle relaxation, vasodilatation, and enhanced penile erection, has been a major advancement in the treatment of erectile dysfunction.⁷ The vasodilating action of sildenafil affects both the arteries and the veins, so the most frequent side effects of sildenafil are headache and facial flushing.⁸ Sildenafil causes small decreases in systolic and diastolic blood pressures, but clinically significant hypotension is rare. Studies of sildenafil and nitrates taken together show much greater drops in blood pressure. For that reason, it is contraindicated to use sildenafil in patients who take long-acting nitrates or who use short-acting, nitrate-containing medications.

In the phase II/III studies completed before Food and Drug Administration (FDA) approval, >3700 patients received sildenafil and almost 2000 received placebo in double-blind and open-label studies. None were taking long-acting nitrates, although patients with coronary artery disease were not excluded. Approximately 25% of the patients had hypertension and were taking antihypertensive medications, and 17% were diabetic. In these studies, the incidence of serious cardiovascular adverse effects was similar in the double-blind sildenafil group, the double-blind placebo group, and the open-label group. There were 28 patients who had a myocardial infarction. When adjusted for patient-years of exposure, there were no differences in myocardial infarction rate between the sildenafil group and the placebo group, and no deaths were attributed to treatment. The incidence of myocardial infarction was 1.7/100 patient-years (95% CI, 0.8 to 2.6) in the sildenafil group and 1.4/100 patient-years (95% CI, 0.2 to 2.6) in the placebo group.⁹ In the subsequent analysis done in May 1998, sildenafil exposure had increased to 4913 patient-years (693 double-blind sildenafil; 4220 open-label extensions), and 26 deaths had been reported, for an incidence rate of 0.53/100 patient-years. The incidence for placebo remained the same (ie, 2 deaths or 0.57/100 patient-years).⁵

There have now been >3.6 million prescriptions¹⁰ written for sildenafil, and 4500 patients taking sildenafil have been followed up without any change in the above conclusions. A total of 69 deaths have been reported to the FDA as of August 26, 1998, in patients who have used Viagra.^{10,11} Twenty-one were due to unknown causes, 2 due to stroke, and 46 related to probable cardiac events.^{10,11} Twelve deaths involved a possible interaction between Viagra and nitrates.^{10,11}

Patients with erectile dysfunction are mostly over age 45 and are in general more likely to have risk factors predisposing them to cardiovascular disease, including myocardial infarction and stroke. The vast majority of patients in the clinical development program did not have known coronary disease or congestive heart failure, nor were hypertensive patients taking complicated, multidrug, antihypertensive medical regimens included in the program. Furthermore, 62% of the patients taking Viagra were within the 45- to 64-year-

old age category, and only 23% were aged ≥ 65 years (Pfizer Inc, unpublished data). Although sildenafil is not presently indicated in women, the cautions referred to in this document should probably apply to both men and women, pending studies performed specifically in women.

B. Development of an ACC Expert Consensus Document

In July 1998, responding to inquiries from both concerned physicians and the press, ACC president Spencer King asked the ACC Technology and Practice Executive Committee (TPEC) to supervise the writing of a press release, summary statement, and Expert Consensus Document on sildenafil (Viagra). This article was written to appropriately caution and not to unduly alarm physicians in their use of sildenafil in patients with heart disease.

Dr King and TPEC chair Dr James Forrester selected a group of physicians with specific expertise to prepare the document. Drs Melvin Cheitlin and Adolph Hutter, Jr, were chosen as cochairs of the Writing Group, on the basis of their status as well-recognized senior clinical cardiologists and their experience in producing clinical practice guidelines. Other members were selected for specific expertise: Dr Brindis (managed care), Dr Ganz (vascular reactivity), Dr Kaul (nitric oxide donors), and Dr Zusman (pharmacology of antihypertensive agents). Dr King also invited the AHA to jointly author the document. Dr Richard Russell (critical care cardiology) was appointed to the Writing Group by AHA president Dr Valentin Fuster. All members of the Writing Group were asked to carefully review any potential conflicts of interest they might have regarding their industry relationships. Those writers who indicated conflicts are identified in the byline.

The Writing Group reviewed both the limited published data on Viagra and unpublished data provided by the manufacturer of Viagra, Pfizer Inc. With respect to the unpublished data, all members of the Writing Group who had access to these documents signed statements that they would not distribute this information outside of the Writing Group until such time as it became public information. Members of the Writing Group were instructed to channel all communications with Pfizer through ACC professional staff to eliminate the appearance of bias.

After completion of the document, 10 external referees reviewed the text. A copy of the draft was also provided to Pfizer and to the FDA for comment. The comments from external review, which were kept anonymous, were provided to the Writing Group, which made revisions as they deemed appropriate. The Expert Consensus Document was approved by vote of the TPEC for presentation to the ACC Board of Trustees, which voted to approve its publication in the *Journal of the American College of Cardiology*. The AHA Scientific Advisory Committee also reviewed and approved this document for publication in *Circulation*.

II. Background

A. Physiology of Erection

Penile erection is accomplished by engorgement of cavernous spaces within the corpora cavernosa under near-arterial pres-

ures and involves dilation of arterial inflow, relaxation of corpora cavernosa smooth muscle, and constriction of venous outflow.¹² The blood flow to the penis is supplied by the cavernosal arteries and their branches, the helicine arteries, which empty directly into the cavernous spaces.¹² Erection is initiated by dilation of helicine arteries, resulting in marked augmentation of blood inflow and transmission of arterial pressures to the cavernosal spaces. Relaxation of smooth muscle trabeculae surrounding cavernosal spaces facilitates blood pooling and engorgement. Restriction of venous outflow is also essential to entrapment of blood in the corpora cavernosa and is caused by compression of venules by the expanding smooth muscle trabeculae against the thick tunica albuginea.¹²

B. Role of Nitric Oxide and cGMP

The relaxation of the penile arterial smooth muscle, the corporal smooth muscle, and therefore erection is under the control of the autonomic nervous system.¹³ The principal neural mediator of penile smooth muscle relaxation is nitric oxide (NO).^{13,14} NO and its derivatives have received much attention because they also account for the biological activity of the endothelium-derived relaxation factor and of organic and inorganic nitrate vasodilators. Three isoforms of NO synthase (NOS) that convert L-arginine to NO have been identified: neuronal (nNOS; type I NOS), inducible (iNOS; type II NOS), and endothelial (eNOS; type III NOS). Terminals containing nNOS densely innervate the corpus cavernosum and its arterial supply.^{13,14} NO derived from the endothelium lining penile arteries and cavernosal sinuses also participates in the erectile response. The arterial dilator actions of NO and its relaxant effect on the smooth muscle of the corpus cavernosum are mediated by the activation of soluble guanylate cyclase and production of cGMP, which acts as a second messenger.^{13,14} Accumulation of cGMP leads to a reduction in intracellular calcium and smooth muscle relaxation. The degradation of cGMP into its inactive form, GMP, is catalyzed by cyclic nucleotide phosphodiesterase enzymes.^{15,16} The predominant isoform of this enzyme in the corpus cavernosum is PDE5.^{12,15} Inhibitors of the activity of this enzyme prevent the breakdown of cGMP, resulting in enhanced penile erection.

III. Sildenafil

A. Introduction and Mechanism of Action

Sildenafil belongs to a class of compounds called PDE inhibitors. PDEs comprise a diverse family of enzymes that hydrolyze cyclic nucleotides (cAMP and cGMP) and therefore play a critical role in the modulation of second-messenger signaling pathways.¹⁵

Sildenafil is a potent and selective inhibitor of cGMP-specific PDE5 (Pfizer, unpublished data), the predominant isozyme that metabolizes cGMP in the corpus cavernosum of the penis. cGMP is the second messenger of NO and a principal mediator of smooth muscle relaxation and vasodilatation in the penis. By inhibiting the hydrolytic breakdown of cGMP, sildenafil prolongs the action of cGMP. This results in augmented smooth muscle relaxation and hence prolongation of the erection. Prior production of cGMP by

NO, released primarily from the nonadrenergic, noncholinergic (nitroxidergic) cavernosal nerves in response to sexual stimulation, is required for sildenafil to be effective.^{13,14}

Relatively high levels of PDE5 are found in the human corpus cavernosum; in vascular, visceral, and tracheal smooth muscle; and in platelets.¹⁵ Sildenafil is a potent inhibitor of PDE5, with favorable selectivity (>1000-fold) for human PDE5 over human PDE2 (isozyme found predominantly in the adrenal cortex),¹⁵ PDE3 (found predominantly in smooth muscles, platelets, and cardiac tissue),¹⁵ and PDE4 (found predominantly in the brain and lung lymphocytes)¹⁵ and moderate selectivity (>80-fold) over PDE1 (a cGMP-hydrolyzing isozyme found predominantly in the brain, kidney, and smooth muscle).¹⁵ Sildenafil is only \approx 10-fold as potent for PDE5 as for PDE6 (an enzyme found in the photoreceptors of the human retina); this lower selectivity is presumed to be the basis for abnormalities related to color vision observed with higher doses or plasma levels of sildenafil (Pfizer, unpublished data). The \approx 4000-fold greater selectivity for PDE5 over PDE3 is important because inhibitors of PDE3 (the isozyme involved in regulation of cardiac contractility), such as milrinone, vesnarinone, and enoximone, that have been used in patients with heart failure are generally associated with increased incidence of cardiac arrhythmias and other serious side effects.¹⁷

B. Pharmacokinetics and Metabolism

Sildenafil is rapidly absorbed after oral administration, with absolute bioavailability of \approx 40%. Plasma concentrations peak within 30 to 120 minutes (median, 60 minutes) of oral dosing in the fasted state. Sildenafil is primarily metabolized by the cytochrome P450 3A4 (major route) and 2C9 (minor route) hepatic microsomal isoenzymes, which convert it to an active *N*-desmethyl metabolite that has been shown to possess 50% of the parent drug's potency for inhibiting PDE5. Plasma concentrations of this metabolite are \approx 40% of those seen for sildenafil, so that the metabolite accounts for \approx 20% of the pharmacological effects of sildenafil. Sildenafil and its active metabolite are both highly bound to plasma proteins (\approx 96%), and their terminal half-lives are \approx 4 hours each. The mean steady-state volume of distribution for sildenafil is 105 L, indicating distribution into the tissues. Sildenafil is excreted as metabolites predominantly in the feces (\approx 80% of administered oral dose) and to a lesser extent in the urine (\approx 13% of the administered oral dose). Less than 0.001% of the administered dose appears in the semen; this dose is very unlikely to have any effects in the partners of patients taking sildenafil. Plasma levels of sildenafil are increased in patients aged >65 years (40% increase) and in patients with hepatic impairment (eg, cirrhosis; 80% increase), severe renal impairment (creatinine clearance <30 mL/min; 100% increase), and concomitant use of potent cytochrome P450 3A4 inhibitors (eg, macrolide antibiotics such as erythromycin [200% increase] and clarithromycin; cimetidine; and antifungal agents such as ketoconazole and itraconazole).¹⁸ Protease inhibitors such as indinavir, zidovudine, zalcitabine, and zalcitabine have not been formally studied but, being potent 3A4 inhibitors, are anticipated to have similar effects on sildenafil metabolism (Pfizer, unpublished data).

C. Pharmacodynamics

The pharmacodynamic end points that have been investigated with sildenafil reflect the distribution of PDE5 in different tissues, ie, human corpus cavernosum (penile tumescence), vascular smooth muscle (vasodilatation), and platelets (anti-platelet function).

1. Effects on Penile Tumescence

The efficacy of sildenafil in enabling patients with erectile dysfunction due to a broad spectrum of causes, including vasculogenic (diabetes), neuroreflexogenic (spinal cord injury), and psychogenic (nonorganic), to achieve and maintain erection sufficient for satisfactory sexual intercourse has been demonstrated in all 21 double-blind, randomized, placebo-controlled, multicenter studies (Pfizer, unpublished data).

2. Cardiovascular Effects

a. Effects on Cardiac Contractility

Unlike cAMP-specific PDE3 inhibitors (milrinone, vesnarinone, and enoximone) that increase long-term mortality in patients with heart failure,^{17,19} sildenafil is highly selective (>4000-fold) for human PDE5 over human PDE3 and has not been found to elevate cAMP (Pfizer, unpublished data). The cardiotoxic effects of PDE3 inhibitors are thought to be related to increases in intracellular cAMP in the myocardium.^{15,19,20} Furthermore, PDE5 is not present in cardiac myocytes, and sildenafil has been shown to have no direct inotropic effects on dog trabeculae muscle (Pfizer, unpublished data). However, sildenafil has not been investigated extensively in heart failure patients.

b. Effects on Blood Pressure and Heart Rate

Sildenafil produces a transient modest reduction in systolic (8 to 10 mm Hg) and diastolic (5 to 6 mm Hg) blood pressures, with peak effects evident at 1 hour after the dose (coincident with peak plasma concentrations) and returning to baseline values by 4 hours after the dose (Pfizer, unpublished data). No significant effects are observed on heart rate. The hypotensive effects of sildenafil are neither age dependent (similar reductions in blood pressure in patients aged <65 years compared with those >65 years) nor dose related (over the range of 25 to 100 mg) and rarely result in reports of orthostatic effects. Doses as high as 800 mg have been well tolerated in some healthy volunteers.¹³

c. Effects on Central Hemodynamics and Peripheral Vasculature

In normal volunteers, no significant changes in cardiac index were evident up to 12 hours after the dose for oral sildenafil (100 to 200 mg) or intravenous sildenafil (20 to 80 mg) (Pfizer, unpublished data). Significant decreases in systemic vascular resistance index were reported at the end of intravenous sildenafil infusion (20 to 80 mg), when plasma concentrations were highest (Pfizer, unpublished data). Sildenafil has both arteriodilator and venodilator effects on the peripheral vasculature (Pfizer, unpublished data). In 8 patients with stable angina, intravenous sildenafil reduced systemic and pulmonary arterial pressures and cardiac output by 8%, 25%, and 7%, respectively, consistent with its mixed arterial (systemic and pulmonary hypotension) and venous

(drop in stroke volume secondary to decreased preload) vasodilator effects.¹⁴

In conclusion, consistent with the anticipated effects resulting from an increase in cGMP levels in vascular smooth muscle, sildenafil possesses vasodilatory properties, which result in mild, generally clinically insignificant decreases in blood pressure when taken alone.

d. Platelet Effects

Sildenafil has no direct effects on platelet function but will modestly potentiate the inhibitory effect of the NO donor sodium nitroprusside on ADP-induced platelet aggregation *ex vivo*, consistent with the requirement for an NO drive for sildenafil to produce its pharmacological effects (Pfizer, unpublished data). No effects on bleeding or prothrombin times were seen in healthy subjects receiving sildenafil alone or concurrently with aspirin or warfarin. In addition, no adverse bleeding episodes have been reported with the use of sildenafil (Pfizer, unpublished data). However, because the effects of sildenafil have not been evaluated in patients with bleeding disorders or in patients taking nonaspirin antiplatelet agents (eg, ticlopidine, clopidogrel, or dipyridamole), caution should be exercised when the drug is administered in these clinical settings.

3. Effects on Visual Function

Transient visual abnormalities (mostly color-tinged [blue-green] vision, increased perception of light, and blurred vision) have been reported in patients taking sildenafil, especially at high oral doses (>100 mg) (Pfizer, unpublished data). These visual effects appear to be related to the weaker inhibiting action of sildenafil on PDE6, which regulates signal transduction pathways in the retinal photoreceptors. Sildenafil is 10-fold selective for PDE5 over PDE6 (Pfizer, unpublished data). In patients with inherited disorders of retinal PDE6, such as retinitis pigmentosa, sildenafil should be administered with extreme caution (Pfizer, unpublished data).

4. Adverse Effects

The adverse effects of sildenafil reflect its pharmacological activity of inhibition of PDE5 in various tissues and can be broadly classified into 4 major adverse reactions:

1. *Vasodilatory effects* resulting in headache (16%), flushing (10%), and rhinitis (4%) (the latter presumably as a result of hyperemia of nasal mucosa where PDE5 is present). Dizziness (2%), hypotension (<2%), and postural hypotension (<2%) have been reported rarely and occur at a similar rate in sildenafil- and placebo-treated patients (Pfizer, unpublished data).
2. *Gastrointestinal effects* resulting in dyspepsia and burning sensation from reflux due to relaxation of lower esophageal sphincter (7%) (Pfizer, unpublished data).
3. *Visual abnormalities* resulting in blue-green color-tinged vision, increased perception of light, and blurred vision (3%), especially at higher doses (Pfizer, unpublished data).
4. *Musculoskeletal effects* resulting in myalgias, especially with multiple daily doses. No treatment-related changes in serum creatine kinase or electromyogram have been

observed, however (Pfizer, unpublished data). There is no obvious pharmacological explanation for this effect.

IV. Drug-Drug Interactions and Concomitant Disease States

A. Interaction With Nitrates

The vasodilator actions of nitrates are profoundly amplified with concomitant use of sildenafil, resulting in major hemodynamic compromise and potentially fatal events (Pfizer, unpublished data). This interaction likely applies to all nitrates and NO donors, irrespective of their predominant hemodynamic site of action (see Appendix A for a list of commonly used nitrates). Sildenafil may also potentiate the hypotensive effects of an inhaled form of nitrate, such as amyl nitrate or nitrite, also known as “poppers,” and therefore is contraindicated. Poppers act by dilating blood vessels, and the concurrent recreational use of poppers and sildenafil could result in sudden and marked lowering of blood pressure, which can be potentially serious or even fatal. This interaction may be even more pronounced in patients taking protease inhibitors concurrently (eg, indinavir [Crixivan], ritonavir [Norvir], nelfinavir [Viracept], or saquinavir [Invirase]).

Dietary sources of nitrites, nitrates, and L-arginine (the substrate from which NO is synthesized) do not contribute to the circulating levels of NO in humans and therefore are unlikely to interact with sildenafil. The anesthetic agent nitrous oxide does not undergo any detectable biotransformation and is eliminated unchanged from the body, mostly via the lungs, usually within minutes of its administration. Because it does not form NO in the human body and does not itself activate guanylate cyclase, there is no contraindication to its use after administration of sildenafil.

It is not known how much time must elapse from the time at which a patient takes sildenafil before a nitrate-containing medication might be given without the marked hypotensive effect being produced. On the basis of the pharmacokinetic profile of sildenafil, it can be assumed that the coadministration of a nitrate within the first 24 hours is likely to produce an exaggerated hypotensive response and is therefore contraindicated unless the benefits are determined to far outweigh the risks. After 24 hours, the administration of a nitrate may be considered, but once again, the response to initial dosages must be monitored carefully. In patients in whom the half-life of sildenafil may be prolonged (see below), a more extended period of time from sildenafil administration to nitrate administration may be required. The preferred form of nitrate therapy in this setting would be short-acting intravenous nitroglycerin infusion under close hemodynamic monitoring.

Similarly, all patients taking either sildenafil or nitrates must be warned of the contraindications and the potential consequences of taking sildenafil in the 24-hour interval after taking a nitrate preparation, including sublingual nitroglycerin. Although sublingual nitroglycerin is very short-acting, its need in the previous 24 hours suggests that it may be needed again after sildenafil-enhanced sexual relations. Furthermore, the presence of even trace amounts of nitrates may have unknown effects in combination with sildenafil. The admin-

istration of sildenafil to a patient who has taken a nitrate in the preceding 24 hours is contraindicated.

Appendix A is a listing of nitrate preparations available in the United States. Other preparations may be available in other countries. A careful history of the medications taken by a patient who has taken sildenafil is essential before treatment of the patient for presumed myocardial ischemia or infarction is initiated.

B. Interaction With Antiplatelet Agents

A clinical trial combining sildenafil with aspirin showed no pharmacokinetic interaction between the 2 medications and no additional effect of sildenafil on bleeding time. Dipyridamole is believed to exert antiplatelet effects by at least 2 mechanisms. Its nonspecific PDE action increases platelet cAMP, and it increases plasma adenosine by blocking its reuptake by erythrocytes.²¹ Ticlopidine and clopidogrel produce antiplatelet aggregatory activity by inhibiting ADP-mediated platelet activation.²² No specific interaction studies have been conducted between sildenafil and dipyridamole, ticlopidine, or clopidogrel.

C. Interaction With Other PDE Inhibitors

PDEs are considered to be major mediators of cross talk between different second-messenger signaling pathways,¹⁵ eg, cGMP is known to inhibit PDE3, which hydrolyzes cAMP, thereby resulting in enhanced cAMP levels.^{15,20} This increase in cAMP levels can potentially augment cAMP-mediated effects in various tissues where PDE3 is localized, ie, Ca²⁺ current (I_{Ca}) and inotropy in cardiac myocytes,²³ vascular smooth muscle relaxation,²⁴ and platelet inhibition.²⁵ The risk of precipitating a cardiotoxic, hypotensive, or hemorrhagic event secondary to combining sildenafil with specific PDE3 inhibitors (such as milrinone, vesnarinone, or enoximone) or with nonspecific PDE inhibitors (such as theophylline, dipyridamole, papaverine, and pentoxifylline) is currently unknown, but such effects are unlikely.¹⁷

D. Drug-Drug Interactions Affecting Metabolic Clearance of Sildenafil

Sildenafil is an inhibitor of the cytochrome P450 2C9 metabolic pathway. It is possible that the administration of sildenafil could result in a significant increase in the plasma concentrations of other drugs metabolized through this pathway. Although tolbutamide and warfarin are metabolized by the P450 2C9 pathway, there is no evidence to date that the concomitant administration of sildenafil affects the metabolic clearance of these 2 drugs.

Sildenafil is predominantly metabolized by both the P450 2C9 pathway and the P450 3A4 pathway (a low-affinity but high-capacity system). Thus, potent inhibitors of the P450 3A4 pathway may increase the plasma concentrations of sildenafil and therefore its pharmacological effect. Cimetidine and erythromycin are commonly prescribed drugs that inhibit the P450 3A4 pathway. As indicated in the approved product labeling, the simultaneous administration of either of these agents significantly increases the plasma concentrations of sildenafil; a lower initial dose (25 mg) should be consid-

ered in the coadministration of sildenafil to patients receiving either of these agents.

Many drugs are metabolized by the P450 3A4 pathway but are not inhibitors of the pathway. The coadministration of 1 of these drugs may lead to a competitive inhibition of the metabolism of sildenafil, although the 3A4 system is a high-capacity enzymatic system. The effects of these agents on the half-life, physiological effects, and side effects of sildenafil are unknown; physicians should be cognizant of the potential interaction of such agents. Appendix B includes a partial listing of commonly prescribed drugs metabolized via the P450 3A4 pathway.

E. Concomitant Administration of Antihypertensive Drugs

Sildenafil administration has been associated with reductions in blood pressure (compared with placebo) of as much as 8/5 mm Hg (systolic/diastolic values). In a drug interaction study of sildenafil and amlodipine, the additional blood pressure reduction in the patient population receiving both sildenafil and amlodipine was not significantly different from the population receiving sildenafil and a placebo (Pfizer, unpublished data). Although formal drug-drug interaction studies have not been conducted with the following medications, no increase in blood pressure–related adverse events or systematic enhancement of the blood pressure–lowering effects of thiazide, loop and potassium-sparing diuretics, ACE inhibitors, calcium channel blockers, or α - or β -adrenergic receptor antagonists have been observed in clinical trials. However, the potential for a hypotensive reaction in patients taking antihypertensive medications as well as sildenafil must be considered and the patient alerted to this possibility. Although not supported by data from the clinical trials, there may be a theoretical concern in a patient receiving multiple medications that include antihypertensive therapy and an inhibitor of the metabolic pathway (cytochrome P450 3A4) of sildenafil.

F. Concomitant Disease States

1. Renal Dysfunction

Patients with severe renal impairment (creatinine clearance <30 mL/min) have a reduced clearance of sildenafil. Plasma levels of the parent drug and of its metabolites in patients with severe renal impairment are approximately twice those found in healthy subjects. Thus, the duration of the effect of sildenafil in these patients will be prolonged and also may be enhanced at any given dosage of the medication. Particular care should be taken in the administration of concomitant medications that may lower blood pressure in patients receiving sildenafil whose renal function is severely impaired. The effects of less-severe degrees of renal dysfunction on the metabolism of sildenafil have been evaluated. There were no significant effects on the metabolism of sildenafil seen in subjects with mild (creatinine clearance 50 to 80 mL/min) or moderate (creatinine clearance 30 to 49 mL/min) renal impairment.²⁴ Of note, the plasma creatinine concentration of the elderly patient with a lower body mass may not accurately reflect the patient's creatinine clearance, and thus initiation of

therapy at 25 mg rather than 50 mg may be appropriate in the elderly.

2. Hepatic Dysfunction

Patients with hepatic dysfunction have a decreased clearance of sildenafil compared with normal subjects. Plasma concentrations of sildenafil and of its metabolites may be significantly increased in patients with hepatic dysfunction. Under such conditions, the duration of activity of sildenafil may be prolonged and the extent of its effects enhanced. As in patients with renal dysfunction, the initiation of therapy at 25 mg rather than 50 mg may be appropriate in patients with hepatic dysfunction.

V. Cardiovascular Effects of Sexual Intercourse in Patients With Coronary Artery Disease

There is potential for a high incidence of overt and covert coronary artery disease in patients with erectile dysfunction on the basis of the epidemiological profiles of both patient groups. Therefore, when prescribing sildenafil, physicians should consider the potential implications of coronary artery disease in sedentary patients who plan to resume sexual activity. Because nitrates are contraindicated for the management of coronary ischemic syndromes in patients taking sildenafil, review of the patient's ability to tolerate the cardiovascular stresses involved with sexual intercourse, particularly patients with coronary artery disease or at increased risk of coronary artery disease, may aid the treating physician in patient management.

Cardiac and metabolic expenditures during sexual intercourse will vary depending on the type of sexual activity. In a laboratory setting, healthy males with their usual female partners achieved an average peak heart rate of 110 bpm with woman-on-top coitus and an average peak heart rate of 127 bpm with man-on-top coitus.²⁶ When oxygen uptake was measured in these men, an average metabolic expenditure during stimulation and orgasm of 2.5 metabolic equivalents (METs) for woman-on-top coitus and 3.3 METs for man-on-top coitus was attained. There was a significant individual variation of cardiovascular responses among patients ranging from 2.0 to 5.4 METs for man-on-top coitus. Thus, to simply equate a level of cardiac or metabolic expenditure during sexual intercourse to an activity such as "climbing 1 or 2 flights of stairs" may underestimate the level of cardiovascular response in individual patients.

In patients with known coronary artery disease whose antianginal medicines were stopped for study purposes,²⁷ Drory et al compared the electrocardiographic monitoring findings in sexual activity with a near-maximal exercise treadmill test (ETT). Most patients had previous myocardial infarctions and were in New York Heart Association functional class I or II. ECG criteria for ischemia during intercourse were found in one third of the patients; two thirds of the time, this was silent rather than symptomatic ischemia. All patients with ischemia during coitus also demonstrated ischemia at ETT. Drory et al also noted significant variation in heart rate response to coitus, with an average heart rate of 118 bpm but with some patients attaining a heart rate of 185

bpm at orgasm. Other small studies with ECG monitoring during intercourse in patients with coronary artery disease concluded that sexual activity may provoke increased ventricular ectopic activity that is not necessarily elicited by other stimuli.²⁸ Jackson²⁹ found that in 19 patients with ischemic heart disease who developed angina during sexual intercourse, these symptoms were abolished with β -blockade. The mean maximum heart rate during sexual intercourse with and without use of β -blockers was 82 and 122 bpm, respectively. This would suggest that these patients may have different hemodynamics while taking antianginal medication that may afford them some protection or lower their risk of ischemia. It should be emphasized that coital death is rare, encompassing only 0.6% of sudden death cases.³⁰ Muller et al³¹ found by retrospective case-crossover methodology that although sexual activity can trigger the onset of myocardial infarction, the relative risk in the 2 hours after sexual activity is very low (2.5; 95% CI, 1.7 to 3.7). Furthermore, sexual activity was a likely contributor to the onset of myocardial infarction only 0.9% of the time. Additionally, they found that the relative risk of myocardial infarction is not increased in patients with a prior history of cardiac disease and that regular exercise appears to prevent triggering. It should be cautioned that these reassuring data should not be extrapolated to patients taking sildenafil if they perform at higher cardiac and metabolic expenditures during coitus. The hemodynamic changes associated with sexual activity may be far greater with an unfamiliar than with a familiar partner, in unfamiliar settings, and after excessive eating and consumption of alcohol. The person most at risk is usually middle-aged and having extramarital relations.

The ETT can gauge the potential cardiac stress of sexual activity. If a patient can achieve 5 or 6 METs on the ETT without demonstrating arrhythmias or ischemia electrocardiographically, they most likely are not at high risk for developing myocardial ischemia as a result of their normal sexual activities.

VI. Recommendations for Sildenafil and the Cardiac Patient

A. Prescribing Sildenafil to Patients at Clinical Risk

1. Sildenafil is absolutely contraindicated in patients undergoing any long-acting nitrate drug therapy or using short-acting nitrates because of the risk of developing potentially life-threatening hypotension.
2. If a patient has stable coronary disease, is not taking a long-acting nitrate, has short-acting nitrate use as the only contraindication to sildenafil, and does not appear to need the nitrate on a consistent basis, the physician and the patient should carefully weigh the risks and benefits of sildenafil treatment. If the patient requires nitrates for mild or moderate exercise limitation, sildenafil should probably not be used.
3. All patients taking organic nitrates, even if they have not asked for Viagra, should be informed about the nitrate-sildenafil hypotensive interaction. There is a substantial potential for patients to obtain Viagra from another physician, a friend, or through the "black

market," circumventing healthcare providers who could offer appropriate caution. Because sildenafil also potentiates the hypotensive effect of an inhaled form of nitrate, such as amyl nitrate or poppers, the concurrent recreational use of poppers and sildenafil could result in sudden and marked hypotensive response that could be serious or fatal. This interaction may be more pronounced in patients taking protease inhibitors concurrently (eg, indinavir, ritonavir, nelfinavir, and saquinavir).

4. Similarly, patients must be warned of the contraindication of taking sildenafil in the 24-hour time interval after taking a nitrate preparation, including sublingual nitroglycerin. The administration of sildenafil to a patient who has taken a nitrate in any form in the preceding 24 hours is contraindicated.
5. Although firm data are lacking, pre-Viagra treadmill tests to assess for the presence of stress-induced ischemia in patients with overt and covert coronary artery disease can guide the patient and physician relative to the risk of cardiac ischemia during sexual intercourse. If the patient can achieve ≥ 5 to 6 METS on an ETT without demonstrating ischemia, the risk of ischemia during coitus with a familiar partner, in familiar settings, without the added stress of a heavy meal or alcohol ingestion, is probably low. We wish to stress that the physical and emotional stresses of sexual intercourse can be excessive in some people, particularly those who have not performed this activity in some time and who are not in good condition. These stresses themselves may produce acute ischemia or precipitate myocardial infarction. Such patients should be advised to use common sense and to moderate their physical exertion and their emotional expectations as they begin their experience with taking Viagra.
6. If patients are taking a combination of antihypertensive medications, they should be cautioned about the possibility of sildenafil-induced hypotension. Because both venous and arterial vasodilatation occur with sildenafil, initial monitoring of the blood pressure with the institution of Viagra use would identify patients with an undesired hypotensive blood pressure response. This is an area of particular concern for the patient with congestive heart failure who has a borderline low blood volume and a low blood pressure status as well as for the patient who is following a complicated, multidrug, antihypertensive therapy regimen.

B. Management of Acute Ischemic Syndromes With Patients Taking Sildenafil

1. The physician should try to establish the time of the last dose of sildenafil. Definitive evidence is currently lacking, but it is possible that a precipitous reduction in blood pressure may occur over the initial 24 hours after a dose of sildenafil. Administration of nitrates in this time interval should be avoided. In the event that nitrates are given after sildenafil administration, it is essential to have the capability to support the patient with fluid resuscitation and α -adrenergic agonists if needed. After 24 hours, the administration of a nitrate may be considered, but once again, appropriate caution with careful monitoring of initial dosages must be used. In patients in whom the half-life of sildenafil may be

prolonged, such as in renal and hepatic dysfunction or patients concurrently taking a potent CYP 3A4 inhibitor, a more extended period of time from sildenafil administration to the time of nitrate administration may be required. In patients with recurring mild angina after sildenafil use, other nonnitrate antianginal agents, such as β -blockers, should be considered.

2. Patients taking sildenafil who have an acute myocardial infarction should be treated in the usual manner as described in the ACC/AHA clinical practice guidelines³² including, where appropriate, primary angioplasty or thrombolytics. The only difference is that nitrates are contraindicated for these patients. If the patient had already used nitrates and sildenafil together, the acute myocardial infarction may have been caused by the low diastolic perfusion pressure of the coronary circulation. Blood pressure support may be sufficient to prevent further myocardial damage if no acute plaque rupture is present.
3. In patients with unstable angina, therapy should include only nonnitrate antianginal medications but should otherwise adhere to principles established in the clinical practice guideline available from the Agency for Health Care Policy and Research.³³ To date, there is no evidence of significant interactions between sildenafil and heparin, β -adrenergic blockers, calcium channel blockers, narcotics, or aspirin. These agents can be used as appropriate. After 24 hours, nitrates may be administered if close monitoring is provided and proper facilities are available for fluid and vasopressor support.

C. Treatment of the Hypotensive Patient With Inadvertent Sildenafil-Nitrate Combination Effect

In patients who inadvertently received nitrates while taking sildenafil and who manifest a severe hypotensive response, nitrate and nitroprusside (ie, NO donor) therapy should be immediately stopped. Depending on clinical circumstances, any of the following therapies should be considered alone or in combination:

1. Place the patient in Trendelenburg position.
2. Provide aggressive fluid resuscitation.
3. Provide judicious use of an intravenous α -adrenergic agonist such as phenylephrine (Neo-Synephrine).
4. Provide an α - and β -adrenergic agonist (norepinephrine) for blood pressure support, with the realization that this could exacerbate or lead to an acute ischemic syndrome.
5. Provide intra-aortic balloon counterpulsation.

D. Limitations and Unresolved Issues

Expert Consensus Documents, as noted in the preamble, are often written in circumstances in which the evidence base and experience with the technology or practice are limited. This is clearly the case with Viagra. The evidence base had significant limitations, and many important issues remain unresolved. Of special significance to the current report is the fact that the preapproval clinical trials of Viagra excluded certain high-risk groups of patients with significant cardiac disease (ie, patients with heart failure, patients with myocardial infarction or stroke within 6 months, or patients with uncontrolled hypertension) or patients with blood pressures of

<90/50 or >170/100 mm Hg. More research needs to be done to assess the specific risks of Viagra use among these cardiovascular patients.

The authors of this Expert Consensus Document identified a number of other unresolved issues that could affect clinical management of the cardiovascular consequences of sildenafil use, including the following:

1. Interaction with nonaspirin antiplatelet agents (eg, ticlopidine, clopidogrel, and dipyridamole).
2. Interaction with other PDE inhibitors, including specific PDE inhibitors (eg, milrinone, vesnarinone, and enoximone) and nonspecific PDE inhibitors (eg, theophylline, dipyridamole, papaverine, and pentoxifylline).
3. Central nervous system effects of sildenafil (PDE5 is present in the brain).
4. Hypotensive effects with sildenafil alone in high-risk cardiac patients (severe heart failure).
5. Musculoskeletal effects (myalgias with chest pains that could be confused with angina).

As more evidence is accumulated, the ACC will consider an update of this Expert Consensus Document.

Appendix A

List of Representative Organic Nitrates

Nitroglycerin

Deponit
Minitran
Nitrok
Nitro-Bid
Nitrocine
Nitroderm
Nitro Disc
Nitro-Dur
Nitrogard
Nitroglycerin
Nitroglycerin T/R
Nitroglyn
Nitrol ointment
Nitrolingual spray
Nitrong
Nitro-Par
Nitropress
Nitro SA
Nitrospan
Nitrostat
Nitro-trans system
Nitro transdermal
Nitro-Time
Transiderm-Nitro
Tridil

Isosorbide Mononitrate

Imdur
ISMO
Isosorbide mononitrate
Monoket

Isosorbide Nitrate

Dilatrate-SR
Iso-Bid
Isordil
Isordil tembids
Isosorbide dinitrate
Isosorbide dinitrate LA

Sorbitrate
Sorbitrate SA

Pentaerythritol Tetranitrate

Peritrate
Peritrate SA

Erythryl Tetranitrate

Cardilate

Isosorbide Dinitrate/Phenobarbital

Isordil w/PB

Illicit Substances Containing Organic Nitrates

Amyl nitrate or nitrite (It is known that amyl nitrate or nitrite is sometimes abused. In abuse situations, amyl nitrate or nitrite may be known by various names, including "poppers.")

Appendix B

Drugs That Are Metabolized by or That Inhibit Cytochrome P450 3A4

Antibiotic/Antifungal

Biaxin (clarithromycin)
Clotrimazole
Erythromycin
Diflucan
Sporanox
Ketoconazole
Miconazole
Noroxin
Troleandomycin

Cardiovascular

Amiodarone
Norvast
Digitoxin
Diltiazem
Disopyramide
Plendil (felodipine)
DynaCirc (isradipine)
Cozaar (losartan)
Posicor (mibefradil)
Nifedipine
Quinidine
Verapamil

HMG

Lipitor (atorvastatin)
Baycol (cerivastatin)
Mevacor (lovastatin)
Zocor (simvastatin)

Central Nervous System

Alprazolam
Carbamazepine
Prozac (fluoxetine)
Luvox (fluvoxamine)
Imipramine
Serzone (nefazodone)
Phenobarbital
Phenytoin
Zoloft
Triazolam

Other

Acetaminophen
Hismanal (astemizole)
Tagamet (cimetidine)
Propulsid (cisapride)
Cyclosporine
Dexamethasone

Ethinyl estradiol
 Naringenin (grapefruit juice)
 Prilosec (omeprazole)
 Rifampin
 Tacrolimus
 Seldane (terfenadine)
 Theophylline
 Rezulin (troglitazone)
 Viagra (sildenafil)
 Protease inhibitors: Crixivan (indinavir), Norvir (ritonavir), Viracept (nelfinavir), Invirase (saquinavir)

Biaxin is a registered trademark of Abbott Laboratories. Diflucan, Norvast, and Zoloft are registered trademarks of Pfizer Inc. Sporanox, Hismanal, and Propulsid are registered trademarks of Janssen Pharmaceutica Inc. Noroxin, Cozaar, Mevacor, and Zocor are registered trademarks of Merck & Co, Inc. Plendil and Prilosec are registered trademarks of Astra Merck Inc. DynaCirc is a registered trademark of Norartis Pharmaceuticals Corporation. Posicor is a registered trademark of Roche Pharmaceuticals. Lipitor and Rezulin are registered trademarks of Parke-Davis. Baycol is a trademark of Bayer Corporation. Prozac is a registered trademark of Eli Lilly and Company. Luvox is a registered trademark of Solvay Pharmaceuticals, Inc. Serzone is a registered trademark of Bristol-Myers Squibb Company. Tagamet is a registered trademark of SmithKline Beecham Pharmaceuticals. Seldane is a registered trademark of Hoechst Marion Roussel. Viagra is a trademark of Pfizer Inc.

Staff, American College of Cardiology

Christine W. McEntee, Executive Vice President
 David P. Bodycombe, ScD, Senior Researcher, Scientific and Research Services
 Kristi R. Mitchell, MPH, Researcher, Scientific and Research Services
 Renee L. Hemsley, Document Development and Practice Guidelines
 Gwen C. Pigman, MLS, Assistant Director, On-Line and Library Services

References

1. NIH Consensus Conference. Impotence: NIH Consensus Development Panel on Impotence. *JAMA*. 1993;270:83-90.
2. Shabsigh R, Fishman IJ, Scott FB. Evaluation of erectile impotence. *Urology*. 1988;32:83-90.
3. Furlow WL. Prevalence of impotence in the United States. *Med Aspects Hum Sex*. 1985;19:13-16.
4. Tardif GS. Sexual activity after a myocardial infarction. *Arch Phys Med Rehabil*. 1989;70:763-766.
5. Mitka M. Viagra leads as rivals are moving up. *JAMA*. 1998;280:119-120. News.
6. American Heart Association. 1998 *Heart and Stroke Statistical Update*. Dallas, Tex: American Heart Association; 1997.
7. Boolell M, Gepi-Attee S, Gingell JC, Allen MJ. Sildenafil, a novel effective oral therapy for male erectile dysfunction [see comments]. *Br J Urol*. 1996;78:257-261.
8. Goldstein I, Lue TF, Padma-Nathan H, Rosen RC, Steers WD, Wicker PA. Oral sildenafil in the treatment of erectile dysfunction. *N Engl J Med*. 1998;338:1397-1404.
9. Morales A, Gingell C, Collins M, Wicker PA, Osterloh IH. Clinical safety of oral sildenafil citrate (VIAGRA) in the treatment of erectile dysfunction. *Int J Impot Res*. 1998;10:69-74.
10. *Summary of Death Reports in Viagra Received from Marketing (late March) through June 1998*. Washington, DC: Food and Drug Administration (Center for Drug Evaluation and Research); 1998.
11. *Viagra Deaths of Symptoms Leading to Death Occurring Within Four to Five Hours*. New York, NY: Pfizer Inc; 1998. Internal document.
12. Andersson KE, Wagner G. Physiology of penile erection. *Physiol Rev*. 1995;75:191-236.

13. Naylor AM. Endogenous neurotransmitters mediating penile erection. *Br J Urol*. 1998;81:424-431.
14. Rajfer J, Aronson WJ, Bush PA, Dory FJ, Ignarro LJ. Nitric oxide as a mediator of relaxation of the corpus cavernosum in response to nonadrenergic, noncholinergic neurotransmission. *N Engl J Med*. 1992;326:90-94.
15. Beavo JA. Cyclic nucleotide phosphodiesterases: functional implications of multiple isoforms. *Physiol Rev*. 1995;75:725-748.
16. McDonald LJ, Murad F. Nitric oxide and cyclic GMP signaling. *Proc Soc Exp Biol Med*. 1996;211:1-6.
17. Packer M, Carver JR, Rodeheffer RJ, et al. Effect of oral milrinone on mortality in severe chronic heart failure: the PROMISE Study Research Group [see comments]. *N Engl J Med*. 1991;325:1468-1475.
18. Terrett NK, Bell AS, Brown D, et al. A potent and selective inhibitor of type 5 cGMP phosphodiesterase with utility for the treatment of male erectile dysfunction. *Bioorg Med Chem Lett*. 1996;1819-1824.
19. Nony P, Boissel JP, Lievre M, et al. Evaluation of the effect of phosphodiesterase inhibitors on mortality in chronic heart failure patients: a meta-analysis. *Eur J Clin Pharmacol*. 1994;46:191-196.
20. Degerman E, Belfrage P, Manganiello VC. Structure, localization, and regulation of cGMP-inhibited phosphodiesterase (PDE3). *J Biol Chem*. 1997;272:6823-6826.
21. Saliel E, Ward A. Ticlopidine: a review of its pharmacodynamic and pharmacokinetic properties, and therapeutic efficacy in platelet-dependent disease states. *Drugs*. 1987;34:222-262.
22. Schror K. The effect of prostaglandins and thromboxane A2 on coronary vessel tone: mechanisms of action and therapeutic implications. *Eur Heart J*. 1993;suppl 1:34-41.
23. Mery PF, Pavoine C, Belhassen L, Pecker F, Fischmeister R. Nitric oxide regulates cardiac Ca²⁺ current: involvement of cGMP-inhibited and cGMP-stimulated phosphodiesterases through guanylyl cyclase activation. *J Biol Chem*. 1993;268:26286-26295.
24. Luginier C, Komars N. Modulation of vascular cyclic nucleotide phosphodiesterases by cyclic GMP: role in vasodilatation. *Eur Heart J*. 1993;14(suppl 1):141-148.
25. Maurice DH, Haslam RJ. Molecular basis of the synergistic inhibition of platelet function by nitrovasodilators and activators of adenylate cyclase: inhibition of cyclic AMP breakdown by cyclic GMP. *Mol Pharmacol*. 1990;37:671-681.
26. Bohlen JG, Held JP, Sanderson MO, Patterson RP. Heart rate, rate-pressure product, and oxygen uptake during four sexual activities. *Arch Intern Med*. 1984;144:1745-1748.
27. Drory Y, Shapira I, Fisman EZ, Pines A. Myocardial ischemia during sexual activity in patients with coronary artery disease [see comments]. *Am J Cardiol*. 1995;75:835-837.
28. Johnston BL, Fletcher GF. Dynamic electrocardiographic recording during sexual activity in recent post-myocardial infarction and revascularization patients. *Am Heart J*. 1979;98:736-741.
29. Jackson G. Sexual intercourse and angina pectoris. *Int Rehabil Med*. 1981;3:35-37.
30. Ueno M. The so-called coition death. *Jpn J Leg Med*. 1963;17:330-340.
31. Muller JE, Mittleman A, Maclure M, Sherwood JB, Tofler GH. Triggering myocardial infarction by sexual activity: low absolute risk and prevention by regular physical exertion: Determinants of Myocardial Infarction Onset study investigators [see comments]. *JAMA*. 1996;275:1405-1409.
32. Ryan TJ, Anderson JL, Antman EM, et al. ACC/AHA guidelines for the management of patients with acute myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Acute Myocardial Infarction). *J Am Coll Cardiol*. 1996;28:1328-1428.
33. Braunwald E, Mark DB, Jones RH, et al. Unstable angina: diagnosis and management. *Clinical Practice Guideline Number 10*. Rockville, Md: Agency for Health Care Policy and Research; 1994. Abstract.

KEY WORDS: sildenafil ■ angina ■ Viagra ■ ACC/AHA Expert Consensus Documents ■ nitric oxide

Use of Sildenafil (Viagra) in Patients With Cardiovascular Disease

Melvin D. Cheitlin, Adolph M. Hutter, Jr, Ralph G. Brindis, Peter Ganz, Sanjay Kaul, Richard O. Russell, Jr, Randall M. Zusman, James S. Forrester, Pamela S. Douglas, David P. Faxon, John D. Fisher, Raymond J. Gibbons, Jonathan L. Halperin, Adolph M. Hutter, Jr, Judith S. Hochman, Sanjiv Kaul, William S. Weintraub, William L. Winters, Jr and Michael J. Wolk
Writing Group Members
Technology and Practice Executive Committee

Circulation. 1999;99:168-177

doi: 10.1161/01.CIR.99.1.168

Circulation is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 1999 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circ.ahajournals.org/content/99/1/168>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation* is online at:
<http://circ.ahajournals.org/subscriptions/>