

Better Population Health Through Behavior Change in Adults A Call to Action

Bonnie Spring, PhD, ABPP, Co-Chair; Judith K. Ockene, Co-Chair, PhD, MEd, MA;
Samuel S. Gidding, MD; Dariush Mozaffarian, MD, DrPH, FAHA; Shirley Moore, RN, PhD;
Milagros C. Rosal, PhD; Michael D. Brown, PhD, FAHA; Dorothea K. Vafiadis, MS, FAHA;
Debbie L. Cohen, MD; Lora E. Burke, PhD, MPH, FAHA; Donald Lloyd-Jones, MD, ScM, FAHA;
on behalf of the American Heart Association Behavior Change Committee of the Council on
Epidemiology and Prevention, Council on Lifestyle and Cardiometabolic Health, Council for High
Blood Pressure Research, and Council on Cardiovascular and Stroke Nursing

The American Heart Association (AHA) has adopted a bold new strategy in framing its 2020 goals: “By 2020, to improve the cardiovascular health of all Americans by 20% while reducing deaths from cardiovascular diseases and stroke by 20%.”¹ By medically treating cardiovascular risk biomarkers and the disease itself, clinicians played a major role in achieving the AHA’s 2010 goal to reduce coronary heart disease, stroke, and risk by 25%.¹ Now, however, with direct annual cardiovascular disease–related costs projected to triple, from \$272 billion in 2010 to \$818 billion in 2030,² economic realities necessitate a new approach. To avoid bankrupting the healthcare system, we must improve the distribution of cardiovascular health levels across the population by preserving cardiovascular health from childhood and by treating health risk behaviors to help more individuals improve their cardiovascular health into older ages. This heightened emphasis on preventing disease by addressing health behaviors leads to 3 novel emphases in the 2020 goals: (1) Preserving positive “cardiovascular health” by promoting healthy lifestyle behaviors; (2) treating unhealthful behaviors (poor-quality diet, excess energy intake, physical inactivity, smoking), in addition to risk biomarkers (adverse blood lipids, high blood pressure, hyperglycemia, obesity); and (3) a combination of individual-level and population-based health promotion strategies that aim to shift the majority of the public toward the next level of improved cardiovascular health.

The 7 metrics that define cardiovascular health (smoking, diet quality, physical activity level, body mass index, blood pressure, blood cholesterol, and fasting blood glucose) are each classified into 3 clinical strata (ideal, intermediate, and poor). Individuals with all 7 metrics at ideal levels are considered to have “ideal cardiovascular health.” However, the prevalence of ideal cardiovascular health is very low in the US population, and the prevalence of poor-quality diet, physical inactivity, and overweight/obesity is alarmingly high, presaging a worsening of the current distribution of health biomarkers.^{1–8} It is abundantly clear that cardiovascular health is being lost from childhood through young adulthood and that the major reasons are adverse health behaviors related to diet, physical activity, healthy weight maintenance, and smoking.⁸ These adverse health behaviors are most prevalent among those with low socioeconomic status, little education, and limited access to health care. The elimination of these health risk behaviors would make it possible to prevent at least 80% of heart disease, stroke, and type 2 diabetes mellitus, and even 40% of cancers.^{1,7,9,10} Given consistent evidence that healthful behavior changes are associated with improved levels of cholesterol, blood pressure, blood glucose, and medical outcomes, health risk behaviors warrant concerted intervention. The AHA’s 2020 goals were designed strategically to help all individuals (from all race and ethnic groups) prevent declines in their current health behaviors and take a step forward toward

The American Heart Association makes every effort to avoid any actual or potential conflicts of interest that may arise as a result of an outside relationship or a personal, professional, or business interest of a member of the writing panel. Specifically, all members of the writing group are required to complete and submit a Disclosure Questionnaire showing all such relationships that might be perceived as real or potential conflicts of interest.

This statement was approved by the American Heart Association Science Advisory and Coordinating Committee on August 14, 2013. A copy of the document is available at <http://my.americanheart.org/statements> by selecting either the “By Topic” link or the “By Publication Date” link. To purchase additional reprints, call 843-216-2533 or e-mail kelle.ramsay@wolterskluwer.com.

The American Heart Association requests that this document be cited as follows: Spring B, Ockene JK, Gidding SS, Mozaffarian D, Moore S, Rosal MC, Brown MD, Vafiadis DK, Cohen DL, Burke LE, Lloyd-Jones D; on behalf of the American Heart Association Behavior Change Committee of the Council on Epidemiology and Prevention, Council on Lifestyle and Cardiometabolic Health, Council for High Blood Pressure Research, and Council on Cardiovascular and Stroke Nursing. Better population health through behavior change in adults: a call to action. *Circulation*. 2013;128:•••–•••.

Expert peer review of AHA Scientific Statements is conducted by the AHA Office of Science Operations. For more on AHA statements and guidelines development, visit <http://my.americanheart.org/statements> and select the “Policies and Development” link.

Permissions: Multiple copies, modification, alteration, enhancement, and/or distribution of this document are not permitted without the express permission of the American Heart Association. Instructions for obtaining permission are located at http://www.heart.org/HEARTORG/General/Copyright-Permission-Guidelines_UCM_300404_Article.jsp. A link to the “Copyright Permissions Request Form” appears on the right side of the page.

(*Circulation*. 2013;128:00-00.)

© 2013 American Heart Association, Inc.

Circulation is available at <http://circ.ahajournals.org>

DOI: 10.1161/01.cir.0000435173.25936.e1

better cardiovascular health by progressing, as needed, from poor to intermediate or from intermediate to ideal healthy lifestyle behaviors.^{5,11} Although full implementation of the goals requires a life-course approach, beginning from childhood, we focus this call to action primarily on needed behavior change for adults.

The Evidence

As evidence that healthy lifestyle change is both necessary and attainable, we highlight the substantial bodies of research that link health behaviors with cardiovascular health and establish that health behavior change is feasible, improves health outcomes, and lowers healthcare costs.

The strong links between individual health behaviors and cardiovascular health were summarized in detail in the AHA's 2020 strategic impact goals statement.¹ Subsequent prospective studies have demonstrated that the presence of ideal cardiovascular health behaviors and biomarkers is associated with longevity and freedom from cardiovascular and all-cause morbidity and mortality. Ford and colleagues⁵ observed that adults who had ≥ 5 of the 7 metrics at ideal levels, compared with 0 metrics, were at 78% lower risk for all-cause mortality and 88% lower risk for cardiovascular disease mortality over 5.8 years of follow-up. Similarly, over 14.8 years of follow-up, Yang and colleagues¹¹ observed a graded reduction in all-cause, cardiovascular disease, and ischemic heart disease mortality in conjunction with having greater numbers of metrics at ideal levels at baseline, which suggests that substantial proportions of deaths could be prevented or postponed with greater achievement of ideal cardiovascular health. Higher numbers of ideal health metrics were also associated with lower cancer mortality.¹¹ Likewise, over 20 years of follow-up, adults in the Atherosclerosis Risk in Communities Study who had higher numbers of either health behaviors or biomarkers at ideal levels had dramatically lower cardiovascular disease event rates, and results were similar for blacks and whites.⁴

Findings from the Coronary Artery Risk Development in Young Adults (CARDIA) study demonstrate that young adults aged 18 to 30 years with healthy behaviors grow up to become older adults with lower biological risk.¹² Among those who practiced all of 5 healthy lifestyle behaviors in young adulthood (nonsmoking, low or no alcohol use, healthy diet, physically active, maintenance of normal weight), 60% had all ideal cardiovascular biomarkers in middle age. In contrast, of those who practiced none of these healthy behaviors in young adulthood, fewer than 5% reached middle age with all cardiovascular biomarkers at ideal levels.¹² Moreover, highlighting the promise of behavior change, those who did not have a healthy lifestyle at baseline but began to practice healthy behaviors later still had higher rates of ideal cardiovascular biomarkers than those who never adopted healthy behaviors. These results indicate that regardless of timing, the adoption of a healthy lifestyle has value for cardiovascular health.¹²

The old folklore that "behavior can't be changed" has been dispelled by numerous randomized controlled trials of effective behavioral interventions.^{13–23} A number of federal agencies' practice guidelines based on this evidence advise clinicians to either provide or refer patients to receive

counseling in health behavior change.^{17–19,24,25} Findings confirm modest but significant impacts of brief physician advice or intervention on smoking cessation, improved dietary quality, and increased physical activity.^{14,15,19,22–27} These small lifestyle changes, as they accrue over time, can yield substantial improvements in the population's cardiovascular disease morbidity and mortality.²⁸ Greater improvement in health behaviors is produced by intensive behavioral treatments, that is, those that incorporate multiple counseling sessions, each lasting ≥ 10 minutes.^{18–20,22–24,29–31} To treat obesity and improve diet quality, the US Preventive Services Task Force endorses and the Patient Protection and Affordable Care Act mandates coverage without copayments only for intensive behavioral intervention, because evidence of effectiveness is insufficient for less intensive interventions.²⁴

Conjoint use of high-risk and population strategies is essential to optimize cardiovascular outcomes and healthcare costs.^{5,32–34} To demonstrate this principle, Milstein and colleagues³⁵ used a dynamic simulation model of the US health system that tested the impact of 3 potential strategies to reduce deaths and lower costs: Expanded health insurance coverage, increased delivery of high-value preventive and chronic care, and promotion of healthier behavior and environments. The model showed that when added to a simulated scenario that included expanded coverage and improved care, health promotion could save 90% more lives and reduce costs by 30% in 10 years. That same investment could, by 25 years, save 140% more lives and reduce costs by 62%.³⁵ In sum, high-risk patients need both medical and intensive behavioral treatments that initially add cost but keep patients from progressing to a state that requires more expensive medical treatments or hospitalization. The promotion of healthier behavior to those in the lower-risk categories through less intensive behavioral intervention and environmental change takes longer to produce a benefit but gradually slows the growth in chronic disease prevalence and lowers costs by reducing the demand for expensive care.

Shifting the population toward healthier behaviors will require time and systems support at the institutional, financial, and policy levels. We call on practitioners to encourage patients toward healthy lifestyle change, while recognizing that many systems changes needed to facilitate provision of such counseling are not under the clinician's control. Consequently, we also call on the healthcare system, insurance companies, employers, and educational institutions to institute policies that align to help shift all sectors of the population toward a healthier lifestyle.

Behavior Change Counseling in Practice

Clinicians play a vital role in fostering healthier behaviors. The fact that $>80\%$ of adults have a usual source of healthcare services gives providers tremendous reach.³⁶ Moreover, most individuals trust their provider's advice, and the patient-provider relationship is longitudinal, which enables follow-up over an extended period of time.²⁴

The 5 A's is a comprehensive, unified, validated treatment algorithm to facilitate health behavior change counseling efficiently within the constraints of the medical visit.^{13,18,19,22} The framework delineates 5 counseling steps that a provider

can complete in several minutes: (1) **Assess** the risk behavior, (2) **advise** change, (3) **agree** on goals and an action plan via shared decision making, (4) **assist** with treatment, and (5) **arrange** follow-up.³⁷ A 5 A's approach has been shown to produce significant changes in a variety of health behaviors, including smoking cessation, dietary change, and physical activity.^{13,26,38-44} However, despite evidence of effectiveness and federal guidelines that recommend use of the 5 A's, behavioral counseling is not delivered to patients consistently.^{21,22} Limited visit time, competing medical priorities, and insufficient financial incentives for health promotion all create barriers.⁴⁵ However, progress has been made, and for both obesity and smoking, a majority of providers now perform the first 2 A's: Assess the risk behavior and advise behavior change.⁴⁶⁻⁴⁸ In contrast, only a small minority continue on to agree on goals, assist with treatment, and arrange follow-up,^{22,47,48} and it is these latter, less frequently performed 3 A's that have the greatest impact on healthful behavior change.^{22,47-51}

Many providers say they omit the last 3 A's because they perceive them as time consuming and feel they lack the needed counseling skills.^{52,53} At the heart of effective counseling is a patient-centered approach, whereby providers collaborate and partner with patients, helping them create plans to reach their own goals.^{26,54,55} Motivational interviewing, originally developed to treat alcohol abuse, is a specific form of patient-centered counseling that has also been applied successfully to foster health behavior changes.⁵⁴⁻⁵⁷ Even without a formal motivational interview, though, providers' use of patient-centered communication strategies improves individuals' confidence and success in changing health behaviors.^{26,41,51} The core counseling strategy is to avoid commanding language and instead to ask open-ended questions and use reflection to express empathy (taking an active interest in the patient's perspective). The clinician first learns what action steps an individual is willing to take and then helps the patient to develop a behavior change plan. By partnering with patients, the provider supports them in assuming the responsibility to improve their own health.

Although they represent a brief, low-intensity approach, the "assist" and "arrange" steps are particularly impactful because they connect patients with the intensive intervention that many need to change health risk behaviors. Intensive behavioral treatments that include more sessions and more active components are well established to produce greater behavior change than interventions with lower intensity.^{21,22} Weight loss and weight loss maintenance, physical activity, diet quality, and smoking cessation outcomes all improve with added treatment sessions, and smoking cessation improves further with the addition of

medications to assist in quitting.^{21,22} Modification of these unhealthy habits challenges patients to acquire skills in self-monitoring, use of behavioral reinforcers, and problem solving, for which expert and peer support are especially beneficial.

Many physicians find that equipping patients with behavioral skills is too tall an order to add to an already jam-packed medical encounter. Fortunately, the movement toward a patient-centered medical home aims to address that challenge. The distribution of care across an integrated, interprofessional team of providers should enable all to work at the top of their training and expertise. Several allied health professions (eg, clinical health psychologists, dietitians, kinesiologists, health educators) have specialized training and expertise in health behavior change. The addition of these experts to the interprofessional healthcare team either by physical collocation, distance consultation, or engagement in staff training can help to achieve higher-quality, more cost-effective care.⁵⁷⁻⁶³

An effort to connect patients with health-promoting influences in their immediate environment also is needed, especially because patients spend ≈ 5000 of their waking hours annually out of contact with any healthcare provider.^{64,65} Support can take the form of helping patients find and connect to neighborhood resources (eg, park district or community center programs, biking trails, farmers' markets). It will undoubtedly also take the form of answering patients' questions about which of an armory of computer programs, applications, sensors, and online communities they should use to support healthy lifestyle changes. Providers will want to become informed about these new technologies, particularly because only a tiny fraction of them have undergone testing for efficacy or harms.⁶⁶⁻⁶⁸ Table 1 lists effective clinician behavior change strategies, as adapted from Artinian et al²² and others.³⁷⁻⁴²

Healthcare Systems to Support Health Behavior Change

Even the most well-intentioned clinician can be thwarted if the healthcare system in place is unhelpful or obstructive. Because providers cannot manage what they do not measure, practical tools are needed to assess health behaviors. Many electronic medical records systems now have fields for tobacco use and body weight, although consistency and accuracy of their use are variable. Assessment of physical activity and dietary behaviors remains more challenging and still traps providers between 2 poor options: Practical but poorly validated brief assessments, or impractical but better-validated long self-report scales and objective measures. The wherewithal to develop more satisfactory alternatives exists,

Table 1. Summary of Effective Clinician Approaches to Help Individuals Adopt Healthier Behaviors

- Implement the 5 A's: (1) **Assess** the risk behavior, (2) **Advise** change, (3) **Agree** on an action plan, (4) **Assist** with treatment, and (5) **Arrange** follow-up
- Adopt a patient-centered approach: Ask open-ended questions; avoid commanding language; help patients create plans to reach their own goals
- Help to increase patients' self-efficacy (confidence) about being able to change behavior by encouraging specific, attainable goals and building on successes
- Ask patients to self-monitor behavior(s) targeted for change by using a paper or Web-based diary or a mobile application
- Encourage use of rewards or incentives that reinforce attainment of behavioral goals
- Advise accessing social support from family, friends, and community programs
- Encourage seeking support from a behavior change specialist as needed
- Schedule regular follow-up (in-person, telephone, and/or electronic) to assess progress

Table 2. Summary of Healthcare Systems Approaches to Support Health Behavior Change

-
- Ongoing training of provider teams on health behavior targets, evidence-based behavior change strategies, and relevant ethnic and cultural tailoring of treatment
 - Practical tools to assess key health behaviors
 - Electronic health record systems that help assess, track, and report on health behaviors for individual patients and entire practices and that link providers to current guidelines
 - Electronic systems that schedule and track initial and follow-up visits for health behavior change
 - Integrated systems and reimbursement policy to support coordinated care by multidisciplinary provider teams (eg, including physicians, nurse practitioners, medical assistants, clinical health psychologists, dietitians, exercise specialists, social workers, and health educators)
 - Restructuring of practice goals and quality benchmarks to incorporate effective dietary, physical activity, tobacco, and adiposity interventions and targets
-

because item response methodology and computer adaptive testing could support the development of brief, psychometrically sound tests that can be administered online.⁶⁹ Moreover, a helpful reimbursement precedent for technology-mediated assessment was established when the Centers for Medicare & Medicaid Services chose to provide reimbursement for a 3-minute mental health screening delivered by *mym3*, a downloadable application.

To date, healthcare systems improvements are more advanced for efforts to control tobacco compared with obesity, poor diet, or physical inactivity. For example, once the US Preventive Services Task Force concluded that provider training about smoking cessation counseling and pharmacological treatment improved the success of patients' attempts to quit smoking,²¹ medical centers developed in-person and online training programs in which providers could earn certification in tobacco treatment. Also, The Joint Commission on Accreditation of Healthcare Organizations first required smoking cessation counseling as a performance measure for pneumonia, heart failure, and myocardial infarction hospital admissions⁷⁰ and then added smoking cessation counseling as a quality criterion for excellence. These measures were successful in increasing counseling,⁷¹ although effects on tobacco use are just beginning to be studied.⁷² Recently, The Joint Commission announced plans for a requirement to assess smoking and offer cessation counseling to all hospitalized patients.⁷³

Similar systems are starting to emerge for obesity. For example, the Centers for Disease Control and Prevention sponsors a Diabetes Training and Technical Assistance Center that offers training in evidence-based intensive treatment for obesity.³¹ In comparison, very little has been done at the healthcare system level to incorporate specific provider training or quality performance measures related to diet or physical activity. Expansion of current healthcare systems approaches that target tobacco control and an aggressive extension of similar systems and strategies that address adiposity, diet quality, and physical activity are urgently needed.

Allied health professionals can extend the medical team's expertise in health behavior change if reimbursement policies are revised to support their effort. They can help train existing members of the medical staff (eg, medical assistants and nurses) in behavior change strategies and can disseminate training to community programs. Tobacco quit lines,⁷⁴ YMCA-based Diabetes Prevention Program treatment,⁶⁰ and peer educator-led Diabetes Prevention Program treatment⁵⁹ are all examples of effective programs that were originally developed by behavior change professionals and then effectively

implemented in the community. Table 2 itemizes specific changes in the healthcare system that can support and sustain clinician-driven efforts to foster healthy lifestyle change.^{63,75–80}

Conclusion and Call to Action

Intervening to achieve sustained improvements in tobacco use, obesity, poor-quality diet, and physical inactivity is feasible and will improve patients' health. Both individual clinical and population-level strategies need to be implemented synergistically to reduce the high prevalence of these risk behaviors in the population (Table 3). The AHA encourages the community of healthcare providers to do 2 things. The first is to intervene directly and as members of an interprofessional healthcare team to help individual patients adopt healthier lifestyles. Providers' efforts to champion a healthy lifestyle as the norm and healthy behaviors as the default in people's lives are critical. The second is to advocate for healthcare system and policy improvements to address the behavior change needs of the entire population more effectively.

An effective healthcare system to improve cardiovascular health by fostering healthy lifestyle change may resemble a 2-tiered, stepped care model.⁸¹ In the first service tier, clinicians provide low-intensity 5 A's intervention to all patients with low and moderate levels of cardiovascular risk. In the second service tier, the interprofessional healthcare team collaborates, drawing in community- and technology-based resources as needed, to provide intensive behavioral intervention to those at moderate and high levels of risk. Support for these endeavors needs to be available from all healthcare professionals and systems, particularly because the goal is to promote healthier lifestyles that reduce risk for all chronic disease.

Table 3. Key Points

-
- Implement individual patient strategies and support population strategies to improve cardiovascular outcomes
 - Assess and track health behaviors, including smoking, weight, diet, and activity, over time electronically
 - Collaborate with an interprofessional healthcare team to help patients adopt healthier lifestyles through delivery of behavior change counseling
 - Engage community resources and technologies to support health behavior change
 - Support reimbursement of intensive behavioral counseling for patients whose poor health habits put them at cardiovascular risk
 - Advocate for healthcare systems and public health policies that prevent the loss of healthful behaviors over the lifespan and that address the behavior change needs of the entire population
-

Disclosures

Writing Group Disclosures

Writing Group Member	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory Board	
							Other	Other
Bonnie Spring	Northwestern University	NIH†; Proteus†	None	None	None	None	Actigraph*; Alere*	None
Judith K. Ockene	University of Massachusetts Medical School	National Cancer Institute†	None	None	None	None	None	None
Michael D. Brown	Temple University	None	None	None	None	None	None	None
Lora E. Burke	University of Pittsburgh	NIH†	None	None	None	None	None	None
Debbie L. Cohen	University of Pennsylvania	NIH†	None	None	Witness for defense for a nephrology case*	None	None	None
Samuel S. Gidding	Nemours Foundation	None	None	None	None	None	None	None
Donald Lloyd-Jones	Northwestern University	None	None	None	None	None	None	None
Dariush Mozaffarian	Brigham and Women's Hospital/Harvard School of Public Health	NIH†; Genes and Environment Initiative at Harvard School of Public Health†; Gates Foundation/World Health Organization†; GlaxoSmithKline; Pronova; Searle Scholar Award from the Searle Funds at the Chicago Community Trust†; Sigma Tau	None	Aramark*; the Chicago Council*; International Life Sciences Institute*; Norwegian Seafood Export Council*; Nutrition Impact*; SPRIM*; Unilever*; Food and Agricultural Organization of the United Nations*; US Food and Drug Administration*; World Health Organization*	None	Harvard has filed a provisional patent application that been assigned to Harvard, listing Dr Mozaffarian as a coinventor for use of trans-palmitoleic acid to prevent and treat insulin resistance, type 2 diabetes mellitus, and related conditions*; royalties from UpToDate for an online chapter*	Foodminds*	None
Shirley Moore	Case Western Reserve University	AHA†; NIH†	None	None	None	None	VA†	None
Milagros C. Rosal	University of Massachusetts Medical School	None	None	None	None	None	None	None
Dorothea K. Vafiadis	American Heart Association	None	None	None	None	None	None	None

This table represents the relationships of writing group members that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all members of the writing group are required to complete and submit. A relationship is considered to be "significant" if (1) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (2) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.

*Modest.

†Significant.

Reviewer Disclosures

Reviewer	Employment	Research Grant	Other Research Support	Speakers' Bureau/Honoraria	Expert Witness	Ownership Interest	Consultant/Advisory	
							Board	Other
Linda J. Ewing	University of Pittsburgh	None	None	None	None	None	None	None
Elizabeth Goodman	Massachusetts General Hospital	None	None	None	None	None	None	None
Kristen Hairston	Wake Forest University	None	None	None	None	None	None	None

This table represents the relationships of reviewers that may be perceived as actual or reasonably perceived conflicts of interest as reported on the Disclosure Questionnaire, which all reviewers are required to complete and submit. A relationship is considered to be "significant" if (1) the person receives \$10 000 or more during any 12-month period, or 5% or more of the person's gross income; or (2) the person owns 5% or more of the voting stock or share of the entity, or owns \$10 000 or more of the fair market value of the entity. A relationship is considered to be "modest" if it is less than "significant" under the preceding definition.


References

- Lloyd-Jones DM, Hong Y, Labarthe D, Mozaffarian D, Appel LJ, Van Horn L, Greenlund K, Daniels S, Nichol G, Tomaselli GF, Arnett DK, Fonarow GC, Ho PM, Lauer MS, Masoudi FA, Robertson RM, Roger V, Schwamm LH, Sorlie P, Yancy CW, Rosamond WD; on behalf of the American Heart Association Strategic Planning Task Force and Statistics Committee. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic impact goal through 2020 and beyond. *Circulation*. 2010;121:586–613.
- Heidenreich PA, Trogon JG, Khavjou OA, Butler J, Dracup K, Ezekowitz MD, Finkelstein EA, Hong Y, Johnston SC, Khera A, Lloyd-Jones DM, Nelson SA, Nichol G, Orenstein D, Wilson PW, Woo YJ; on behalf of the American Heart Association Advocacy Coordinating Committee, Stroke Council, Council on Cardiovascular Radiology and Intervention, Council on Clinical Cardiology, Council on Epidemiology and Prevention, Council on Arteriosclerosis, Thrombosis and Vascular Biology, Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation, Council on Cardiovascular Nursing, Council on the Kidney in Cardiovascular Disease, Council on Cardiovascular Surgery and Anesthesia, and Interdisciplinary Council on Quality of Care and Outcomes Research. Forecasting the future of cardiovascular disease in the United States: a policy statement from the American Heart Association. *Circulation*. 2011;123:933–944.
- Bambs C, Kip KE, Dinga A, Mulukutla SR, Aiyer AN, Reis SE. Low prevalence of "ideal cardiovascular health" in a community-based population: the Heart Strategies Concentrating on Risk Evaluation (Heart SCORE) study. *Circulation*. 2011;123:850–857.
- Folsom AR, Yatsuya H, Nettleton JA, Lutsey PL, Cushman M, Rosamond WD; ARIC Study Investigators. Community prevalence of ideal cardiovascular health, by the American Heart Association definition, and relationship with cardiovascular disease incidence. *J Am Coll Cardiol*. 2011;57:1690–1696.
- Ford ES, Greenlund KJ, Hong YL. Ideal cardiovascular health and mortality from all causes and diseases of the circulatory system among adults in the United States. *Circulation*. 2012;125:987–995.
- Roger VL, Go AS, Lloyd-Jones DM, Adams RJ, Berry JD, Brown TM, Carnethon MR, Dai S, de Simone G, Ford ES, Fox CS, Fullerton HJ, Gillespie C, Greenlund KJ, Hailpern SM, Heit JA, Ho PM, Howard VJ, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Makuc DM, Marcus GM, Marelli A, Matchar DB, McDermott MM, Meigs JB, Moy CS, Mozaffarian D, Mussolino ME, Nichol G, Paynter NP, Rosamond WD, Sorlie PD, Stafford RS, Turan TN, Turner MB, Wong ND, Wylie-Rosett J; on behalf of the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2011 update: a report from the American Heart Association [published corrections appear in *Circulation*. 2011;124:e426 and *Circulation*. 2011;123:e240]. *Circulation*. 2011;123:e18–e209.
- Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, Bravata DM, Dai S, Ford ES, Fox CS, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Makuc DM, Marcus GM, Marelli A, Matchar DB, Moy CS, Mozaffarian D, Mussolino ME, Nichol G, Paynter NP, Soliman EZ, Sorlie PD, Sotoodehnia N, Turan TN, Virani SS, Wong ND, Woo D, Turner MB; on behalf of the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2012 update: a report from the American Heart Association [published correction appears in *Circulation*. 2012;125:e1002]. *Circulation*. 2012;125:e2–e220.
- Shay CM, Ning H, Allen NB, Carnethon MR, Chiuve SE, Greenlund KJ, Davi GL, Lloyd-Jones DM. Status of cardiovascular health in US adults: prevalence estimates from the National Health and Nutrition Examination Surveys (NHANES) 2003–2008. *Circulation*. 2012;125:45–56.
- Institute of Medicine, Committee on Quality of Health Care in America. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington, DC: National Academy Press; 2001.
- Ezzati M, Lopez AD, Rodgers A, Murray CJL. *Comparative Quantification of Health Risks: Global and Regional Burden of Disease Attributable to Selected Major Risk Factors*. Geneva, Switzerland: World Health Organization; 2004.
- Yang Q, Cogswell ME, Flanders WD, Hong Y, Zhang Z, Loustalot F, Gillespie C, Merritt R, Hu FB. Trends in cardiovascular health metrics and associations with all-cause and CVD mortality among US adults. *JAMA*. 2012;307:1273–1283.
- Liu K, Davi GL, Loria CM, Colangelo LA, Spring B, Moller AC, Lloyd-Jones DM. Healthy lifestyle through young adulthood and the presence of low cardiovascular disease risk profile in middle age: the Coronary Artery Risk Development in (Young) Adults (CARDIA) Study. *Circulation*. 2012;125:996–1004.
- Makoul G, Dhurandhar A, Goel MS, Scholtens D, Rubin AS. Communication about behavioral health risks: a study of videotaped encounters in 2 internal medicine practices. *J Gen Intern Med*. 2006;21:698–703.
- Goldstein MG, Whitlock EP, DePue J; Planning Committee of the Addressing Multiple Behavioral Risk Factors in Primary Care Project. Multiple behavioral risk factor interventions in primary care: summary of research evidence. *Am J Prev Med*. 2004;27(suppl):61–79.
- Rees K, Dyakova M, Ward K, Thorogood M, Brunner E. Dietary advice for reducing cardiovascular risk (review). *The Cochrane Library*. March 28, 2013;3:CD002128.
- Stead LF, Bergson G, Lancaster T. Physician advice for smoking cessation. *Cochrane Database Syst Rev*. 2008;(2):CD000165.
- Smith SC Jr, Benjamin EJ, Bonow RO, Braun LT, Creager MA, Franklin BA, Gibbons RJ, Grundy SM, Hiratzka LF, Jones DW, Lloyd-Jones DM, Minissian M, Mosca L, Peterson ED, Sacco RL, Spertus J, Stein JH, Taubert KA. AHA/ACC secondary prevention and risk reduction therapy for patients with coronary and other atherosclerotic vascular disease: 2011 update: a guideline from the American Heart Association and American College of Cardiology Foundation. *Circulation*. 2011;124:2458–2473.
- National Institutes of Health, NHLBI Obesity Education Initiative and North American Association for the Study of Obesity. *The Practical Guide: Identification, Evaluation, and Treatment of Overweight and Obesity in Adults*. Bethesda, MD: National Institutes of Health; October 2000. NIH publication No. 00-4084.
- US Preventive Services Task Force Web site. Topic page. Behavioral counseling in primary care to promote a healthy diet in adults at increased risk for cardiovascular disease. 2003. <http://www.uspreventiveservicestaskforce.org/uspstf/uspstf/diet.htm>. Accessed February 12, 2012.
- Spring B, Schneider K, McFadden HG, Vaughn J, Kozak AT, Smith M, Moller AC, Epstein LH, DeMott A, Hedeker D, Siddique J, Lloyd-Jones DM. Multiple behavior changes in diet and activity: a randomized controlled trial using mobile technology. *Arch Intern Med*. 2012;172:789–796.
- Tobacco Use and Dependence Guideline Panel. *Treating Tobacco Use and Dependence: 2008 Update*. Rockville, MD: US Department of Health and Human Services, Public Health Service; 2008.
- Artinian NT, Fletcher GF, Mozaffarian D, Kris-Etherton P, Van Horn L, Lichtenstein AH, Kumanyika S, Kraus WE, Fleg JL, Redeker NS, Meininger JC, Banks J, Stuart-Shor EM, Fletcher BJ, Miller TD, Hughes S, Braun LT, Kopin LA, Berra K, Hayman LL, Ewing LJ, Ades PA, Durstine JL,

- Houston-Miller N, Burke LE; on behalf of the American Heart Association Prevention Committee of the Council on Cardiovascular Nursing. Interventions to promote physical activity and dietary lifestyle changes for cardiovascular risk factor reduction in adults: a scientific statement from the American Heart Association. *Circulation*. 2010;122:406–441.
23. Burke LE, Styn MA, Sereika SM, Conroy MB, Ye L, Glanz K, Sevcik MA, Ewing LJ. Using mHealth technology to enhance self-monitoring for weight loss: a randomized trial. *Am J Prev Med*. 2012;43:20–26.
 24. Ockene JK, Kristeller J, Pbert L, Hebert JR, Luippold R, Goldberg RJ, Landon J, Kalan K. The Physician-Delivered Smoking Intervention Project: can short-term interventions produce long-term effects for a general outpatient population? *Health Psychol*. 1994;13:278–281.
 25. Ockene JK, Kristeller J, Goldberg RJ, Amick TL, Pekow PS, Hosmer D, Quirk M, Kalan K. Increasing the efficacy of physician-delivered smoking intervention: a randomized clinical trial. *J Gen Intern Med*. 1991;6:1–8.
 26. US Preventive Services Task Force. *The Guide to Clinical Preventive Services, 2012: Recommendations of the U.S. Preventive Services Task Force*. Rockville, MD: Agency for Healthcare Research and Quality; 2011. AHRQ publication No. 12-05154.
 27. National Diabetes Prevention Program. Centers for Disease Control and Prevention Web site. <http://www.cdc.gov/diabetes/prevention/>. Accessed September 13, 2012.
 28. Rose G. Sick individuals and sick populations. *Int J Epidemiol*. 1985;14:32–38.
 29. Fiore MC. US Public Health Service clinical practice guideline: treating tobacco use and dependence. *Respir Care*. 2000;45:1200–1262.
 30. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med*. 2002;346:393–403.
 31. Counseling and interventions to prevent tobacco use and tobacco-caused disease in adults and pregnant women. US Preventive Services Task Force Web site. Topic page. 2009. <http://www.uspreventiveservicestaskforce.org/uspstf/uspstbac2.htm>. Accessed February 12, 2012.
 32. Ford ES, Ajani UA, Croft JB, Critchley JA, Labarthe DR, Kottke TE, Giles WH, Capewell S. Explaining the decrease in U.S. deaths from coronary disease, 1980–2000. *N Engl J Med*. 2007;356:2388–2398.
 33. Spring B. *Translational Behavioral Medicine: a pathway to better health*. TBM. 2011;1:1–3.
 34. Spring B, Moller AC, Coons M. Multiple health behaviors: overview and implications. *J Public Health (Oxf)*. 2012;34(suppl 1):i3–i10.
 35. Milstein B, Homer J, Briss P, Burton D, Pechacek T. Why behavioral and environmental interventions are needed to improve health at lower cost. *Health Aff (Millwood)*. 2011;30:823–832.
 36. Davis D, Schoenbaum SC, Collins KS, Teeney K, Hughes DL, Audet A-MJ. *Room for Improvement: Patients Report on the Quality of Their Health Care*. New York, NY: The Commonwealth Fund; 2002.
 37. Ammerman A, Pignone M, Fernandez L, Lohr K, Jacobs AD, Nester C, Orleans T, Pender N, Woolf S, Sutton SF, Lux LJ, Whitener L. *Counseling to Promote a Healthy Diet*. Rockville, MD: Agency for Healthcare Research and Quality; 2002. Systematic Evidence Review No. 18.
 38. Increasing physical activity: a report on recommendations of the Task Force on Community Preventive Services. *MMWR Recomm Rep*. 2001;50:1–14.
 39. Task Force on Community Preventive Services. Recommendations regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am J Prev Med*. 2001;20(suppl):10–15.
 40. Kahn EB, Ramsey LT, Brownson RC, Heath GW, Howze EH, Powell KE, Stone EJ, Rajab MW, Corso P. The effectiveness of interventions to increase physical activity: a systematic review. *Am J Prev Med*. 2002;22(suppl):73–107.
 41. Ockene IS, Hebert JR, Ockene JK, Saperia GM, Stanek E, Nicolosi R, Merriam PA, Hurley TG. Effect of physician-delivered nutrition counseling training and an office-support program on saturated fat intake, weight, and serum lipid measurements in a hyperlipidemic population: Worcester Area Trial for Counseling in Hyperlipidemia (WATCH). *Arch Intern Med*. 1999;159:725–731.
 42. Burke LE, Wang J, Sevcik MA. Self-monitoring in weight loss: a systematic review of the literature. *J Am Diet Assoc*. 2011;111:92–102.
 43. Whitlock EP, Orleans CT, Pender N, Allan J. Evaluating primary care behavioral counseling interventions: an evidence-based approach. *Am J Prev Med*. 2002;22:267–284.
 44. Yarnall KSH, Pollak KI, Østbye T, Krause KM, Michener JL. Primary care: is there enough time for prevention? *Am J Public Health*. 2003;93:635–641.
 45. Chase EC, McMenamin SB, Halpin HA. Medicaid provider delivery of the 5A's for smoking cessation counseling. *Nicotine Tob Res*. 2007;9:1095–1101.
 46. Jay M, Gillespie C, Schlaier S, Sherman S, Kalet A. Physicians' use of the 5As in counseling obese patients: is the quality of counseling associated with patients' motivation and intention to lose weight? *BMC Health Serv Res*. 2010;10:159.
 47. Alexander SC, Cox ME, Boling Turer CL, Lyna P, Østbye T, Tulsy JA, Dolor RJ, Pollak KI. Do the five A's work when physicians counsel about weight loss? *Fam Med*. 2011;43:179–184.
 48. Unrod M, Smith M, Spring B, DePue J, Redd W, Winkel G. Randomized controlled trial of a computer-tailored intervention to increase smoking cessation counseling by primary care physicians. *J Gen Intern Med*. 2007;22:478–484.
 49. Gordon JS, Andrews JA, Crews KM, Payne TJ, Severson HH. The 5A's vs 3A's plus proactive quitline referral in private practice dental offices: preliminary results. *Tob Control*. 2007;16:285–288.
 50. Aveyard P, Begh R, Parsons A, West R. Brief opportunistic smoking cessation interventions: a systematic review and meta-analysis to compare advice to quit and offer of assistance. *Addiction*. 2012;107:1066–1073.
 51. Cox ME, Yancy WS Jr, Coffman CJ, Ostbye T, Tulsy JA, Alexander SC, Brouwer RJ, Dolor RJ, Pollak KI. Effects of counseling techniques on patients' weight-related attitudes and behaviors in a primary care clinic. *Patient Educ Couns*. 2011;85:363–368.
 52. Forman-Hoffman V, Little A, Wahls T. Barriers to obesity management: a pilot study of primary care clinicians. *BMC Fam Pract*. 2006;7:35.
 53. Ruelaz AR, Diefenbach P, Simon B, Lanto A, Arterburn D, Shekelle PG. Perceived barriers to weight management in primary care: perspectives of patients and providers. *J Gen Intern Med*. 2007;22:518–522.
 54. Epstein RM, Franks P, Fiscella K, Shields CG, Meldrum SC, Kravitz RL, Duberstein PR. Measuring patient-centered communication in patient-physician consultations: theoretical and practical issues. *Soc Sci Med*. 2005;61:1516–1528.
 55. Rollnick S, Mason P, Butler C. *Health Behavior Change: A Guide for Practitioners*. New York, NY: Churchill Livingstone; 1999.
 56. Rubak S, Sandbaek A, Lauritzen T, Christensen B. Motivational interviewing: a systematic review and meta-analysis. *Br J Gen Pract*. 2005;55:305–312.
 57. Tulloch H, Fortier M, Hogg W. Physical activity counseling in primary care: who has and who should be counseling? *Patient Educ Couns*. 2006;64:6–20.
 58. Frank A. A multidisciplinary approach to obesity management: the physician's role and team care alternatives. *J Am Diet Assoc*. 1998;98(suppl 2):S44–S48.
 59. Katula JA, Vitolins MZ, Rosenberger EL, Blackwell CS, Morgan TM, Lawlor MS, Goff DC Jr. One-year results of a community-based translation of the Diabetes Prevention Program: Healthy-Living Partnerships to Prevent Diabetes (HELP PD) Project [published correction appears in *Diabetes Care*. 2012;35:455]. *Diabetes Care*. 2011;34:1451–1457.
 60. Ackermann RT, Finch EA, Brizendine E, Zhou HH, Marrero DG. Translating the Diabetes Prevention Program into the community: the DEPLOY pilot study. *Am J Prev Med*. 2008;35:357–363.
 61. Appel LJ, Clark JM, Yeh HC, Wang NY, Coughlin JW, Daumit G, Miller ER 3rd, Dalcin A, Jerome GJ, Geller S, Noronha G, Pozefsky T, Charleston J, Reynolds JB, Durkin N, Rubin RR, Louis TA, Brancati FL. Comparative effectiveness of weight-loss interventions in clinical practice. *N Engl J Med*. 2011;365:1959–1968.
 62. Agency for Healthcare Research and Quality Web site. The patient-centered medical home: strategies to put patients at the center of primary care. http://www.pcmh.ahrq.gov/portal/server.pt/gateway/PTARGS_0_12547_955661_0_0_18/. Accessed September 26, 2013.
 63. Public Policy Committee of the American College of Physicians; Ginsburg JA, Doherty RB, Ralston JF Jr, Senkeeto N, Cooke M, Cutler C, Fleming DA, Freeman BP, Gluckman RA, Liebow M, McLean RM, Musana KA, Nichols PM, Purtle MW, Reynolds PP, Weaver KM, Dale DC, Levine JS, Stubbs JW. Achieving a high-performance health care system with universal access: what the United States can learn from other countries [published correction appears in *Ann Intern Med*. 2008;148:635]. *Ann Intern Med*. 2008;148:55–75.
 64. Spring B. Health decision making: lynchpin of evidence-based practice. *Med Decis Making*. 2008;28:866–874.
 65. Asch DA, Muller RW, Volpp K. Automated hovering: watching over the 5000 hours. *N Engl J Med*. 2012;367:1–3.
 66. Rao G, Burke LE, Spring BJ, Ewing LJ, Turk M, Lichtenstein AH, Cornier MA, Spence JD, Coons M; on behalf of the American Heart Association Obesity Committee of the Council on Nutrition, Physical Activity and Metabolism; Council on Clinical Cardiology; Council on Cardiovascular

- Nursing; Council on the Kidney in Cardiovascular Disease; Stroke Council. New and emerging weight management strategies for busy ambulatory settings: a scientific statement from the American Heart Association. *Circulation*. 2011;124:1182–1203.
67. Spring B, Duncan JM, Janke EA, Kozak AT, McFadden HG, DeMott A, Pictor A, Epstein LH, Siddique J, Pellegrini CA, Buscemi J, Hedeker D. Integrating technology into standard eight loss treatment: a randomized controlled trial. *JAMA Intern Med*. 2013; 173:105–111.
 68. Kaplan RM, Stone AA. Bringing the laboratory and clinic to the community: mobile technologies for health promotion and disease prevention. *Annu Rev Psychol*. 2013;64:471–498.
 69. Cella D, Riley W, Stone A, Rothrock N, Reeve B, Yount S, Amtmann D, Bode R, Buysse D, Choi S, Cook K, Devellis R, DeWalt D, Fries JF, Gershon R, Hahn EA, Lai JS, Pilkonis P, Revicki D, Rose M, Weinfurt K, Hays R; PROMIS Cooperative Group. The Patient-Reported Outcomes Measurement Information System (PROMIS) developed and tested its first wave of adult self-reported health outcome item banks: 2005–2008. *J Clin Epidemiol*. 2010;63:1179–1194.
 70. Balkstra CR, Fields M, Roesler L. Meeting Joint Commission on Accreditation of Healthcare Organizations requirements for tobacco cessation: the St. Joseph's/Candler Health System approach to success. *Crit Care Nurs Clin North Am*. 2006;18:105–111, xiii.
 71. Joint Commission report shows quality gains. *Healthcare Benchmarks Qual Improv*. 2007;14:52–54.
 72. Reeves GR, Wang TY, Reid KJ, Alexander KP, Decker C, Ahmad H, Spertus JA, Peterson ED. Dissociation between hospital performance of the smoking cessation counseling quality metric and cessation outcomes after myocardial infarction. *Arch Intern Med*. 2008;168:2111–2117.
 73. Tobacco treatment. The Joint Commission Web site. 2012. http://www.jointcommission.org/tobacco_treatment/. Accessed July 29, 2012.
 74. Borland R, Segan CJ. The potential of quitlines to increase smoking cessation. *Drug Alcohol Rev*. 2006;25:73–78.
 75. Bodenheimer T. Helping patients improve their health-related behaviors: what system changes do we need? *Dis Manag*. 2005;8:319–330.
 76. Simpson LA, Cooper J. Paying for obesity: a changing landscape. *Pediatrics*. 2009;123(suppl 5):S301–S307.
 77. Quist-Paulsen P. Cessation in the use of tobacco: pharmacologic and non-pharmacologic routines in patients. *Clin Respir J*. 2008;2:4–10.
 78. Davis D, Galbraith R; American College of Chest Physicians Health and Science Policy Committee. Continuing medical education effect on practice performance: effectiveness of continuing medical education: American College of Chest Physicians Evidence-Based Educational Guidelines. *Chest*. 2009;135(suppl):42S–48S.
 79. Mozaffarian D AA, Benowitz NL, Bittner V, Daniels SR, Franch HA JDJ, Kraus WE, Kris-Etherton PM, Krummel DA, Popkin BM, Whitsel LP, Zakai NA; on behalf of the American Heart Association Council on Epidemiology and Prevention, Council on Nutrition, Physical Activity and Metabolism, Council on Clinical Cardiology, Council on Cardiovascular Disease in the Young, Council on the Kidney in Cardiovascular Disease, Council on Peripheral Vascular Disease, and the Advocacy Coordinating Committee. Population approaches to improve diet, physical activity, and smoking habits: a scientific statement from the American Heart Association. *Circulation*. 2012;126:1514–1563.
 80. Pratt M, Sarmiento OL, Montes F, Ogilvie D, Marcus BH, Perez LG, Brownson RC; Lancet Physical Activity Series Working Group. The implications of megatrends in information and communication technology and transportation for changes in global physical activity. *Lancet*. 2012;380:282–293.
 81. Brown RL. Configuring health care for systematic behavioral screening and intervention. *Popul Health Manag*. 2011;14:299–305.

KEY WORDS: AHA Scientific Statements ■ behavior ■ population health ■ prevention



Circulation
JOURNAL OF THE AMERICAN HEART ASSOCIATION

Better Population Health Through Behavior Change in Adults: A Call to Action
Bonnie Spring, Judith K. Ockene, Samuel S. Gidding, Dariush Mozaffarian, Shirley Moore,
Milagros C. Rosal, Michael D. Brown, Dorothea K. Vafiadis, Debbie L. Cohen, Lora E. Burke
and Donald Lloyd-Jones

Circulation. published online October 7, 2013;
Circulation is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Copyright © 2013 American Heart Association, Inc. All rights reserved.
Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circ.ahajournals.org/content/early/2013/10/07/01.cir.0000435173.25936.e1.citation>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation* is online at:
<http://circ.ahajournals.org/subscriptions/>